

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) Publication number:

**0 245 825
A1**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 87106822.7

(22) Date of filing: 11.05.87

(51) Int. Cl.⁴: C07D 231/04 , C07D 231/06 ,
C07D 231/12 , C07D 261/02 ,
C07D 261/04 , C07D 261/06 ,
C07D 261/08 , C07D 275/02 ,
A61K 31/415 , A61K 31/42 ,
A61K 31/425

(30) Priority: 09.05.86 US 861179
22.09.86 US 910692
06.04.87 US 32730

(43) Date of publication of application:
19.11.87 Bulletin 87/47

(84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

(71) Applicant: WARNER-LAMBERT COMPANY
201 Tabor Road
Morris Plains New Jersey 07950(US)

(72) Inventor: Belliotti, Thomas R.
2435 Twin Lakes Apt. TA
Ypsilanti Michigan 48197(US)
Inventor: Connor, David T.
2453 Antietam
Ann Arbor Michigan 48105(US)
Inventor: Flynn, Daniel L.
3647 Northfield Church Road
Ann Arbor Michigan 48105(US)
Inventor: Kostian, Catherine R.
3581 Pheasant Run
Ann Arbor Michigan 48108(US)
Inventor: Nies, Donald E.
2133 Medford 11
Ann Arbor Michigan 48104(US)

(74) Representative: Mansmann, Ivo
c/o Gödecke AG - Patentabteilung Postfach
569 - Mooswaldallee 1-9
D-7800 Freiburg(DE)

BEST AVAILABLE COPY

(54) Novel styryl pyrazoles, isoxazoles and analogs thereof having activity as 5-lipoxygenase inhibitors and pharmaceutical compositions containing them.


(57) The present invention is 3,5-substituted, isoxazoles, pyrazoles, isothiazoles, and analogs thereof having 5-lipoxygenase or cyclooxygenase inhibiting activity or as a sunscreen.

NOVEL STYRYL PYRAZOLES, ISOXAZOLES AND ANALOGS THEREOF HAVING ACTIVITY AS 5-LIPOX-
YGENASE INHIBITORS AND PHARMACEUTICAL COMPOSITIONS CONTAINING THEM.


This is a continuation in part of United States Serial No. 910,692 filed September 22, 1986, which is a continuation in part of United States Serial No. 861,179 filed May 9, 1986.

The present invention is a novel styryl pyrazoles and analogs thereof as well as pharmaceutical compositions containing them.

5 Styryl isoxazole derivatives having cardiovascular activity are known. For example, European Patent Applications No. 34754 and NO. 5192 and German Application No. 2943-405 having Derwent Abstract Nos. 66318 D/37, 84501 B/47 and 34567 D/20, respectively, disclose a compound of the general formula




20 However, the present compounds differ from such references by a completely different side chain from that shown above linked to the phenyl at the 2-position through an ether group in each reference. Further, European Patent Application No. 5186 reviewed by Derwent Abstract No. 844908/47 discloses an intermediate isoxazole of the formula




for which no pharmaceutical utility is disclosed.

35 Pyrazole derivatives having biological, specifically antiinflammatory, activity are found in Belgian Patent Nos. 819,890 and 844,972 abstracted in Derwent Abstract Nos. 20948W/13 and 09405Y/06, respectively. A similar derivative is disclosed in German Patent No. 2920941 of Derwent Abstract No. 86535C/49. However, each of these disclosed pyrazole derivatives requires substituents on adjacent carbons of the pyrazole ring.


Specifically excluded from the compounds of the present invention are the compounds of French Patent No. 2104932 of Derwent Abstract No. 46150T-B useful as hypocholesterolemic, antiinflammatories, analgesics, sedatives, antipyretics, and in some instances diuretics. The compounds of the French Patent generally have the formula



Finally, of lesser interest numerous imidazoles are known having various pharmaceutical activity. For example, U.S. Patent No. 3812111 and British 1,046,248 disclose compounds of general formula




and



respectively wherein the - represents various substituents. However, the imidazoles are compounds differing from the present invention in that the compounds have a different ring system from the pyrazoles.

Thus, the novel compounds that are the present invention provide activity for use in the treatment of diseases in which 5-lipoxygenase enzyme activity contributes to the pathological condition. For example, the use for the present novel compounds, compositions and methods of use is for allergy, asthma, arthritis, skin disorders, such as psoriasis or acne, inflammation, for example, inflammatory bowel disease or pain, and further, also cardiovascular disorders including infarction, angina, arrhythmias, stroke, migraine, atherosclerosis, ulcers and other conditions particularly benefited by cytoprotective activity. An additional property of the present novel compounds now found to provide usefulness, for example, as sun screens, is absorption of ultraviolet light.

The present invention is a novel compound of the formula (I)



I

and pharmaceutically acceptable salts thereof;
wherein

(1) -- is a single or double bond;

(2) R, R₁, and R₂ may be the same or different and are selected from the group consisting of hydrogen, lower alkyl, hydroxy, OR₃ wherein R₃ is lower alkyl, C(O)OR₄ wherein R₄ is hydrogen or lower alkyl, OC(O)R₃ wherein R₃ is independently as defined above, C(O)R₃ wherein R₃ is independently as defined above, NR₆R₇, wherein R₆ and R₇ may be the same or different and are hydrogen or lower alkyl, NHC(O)R₃ wherein R₃ is independently as defined above, NHCHO, NHSO₂R₃ wherein R₃ is independently as defined above, NHCONHR₄ wherein R₄ is independently as defined above, hydroxymethyl, halogen, trifluoromethyl, SR₄ wherein R₄ is independently as defined above, or nitro;

(3) Q is (CH₂)_n wherein n is an integer of zero to four, CH=CH or


wherein R_4 is independently as defined above;

(4) X and Y are (i) N, (ii) NR_5 wherein R_5 is hydrogen, lower alkyl,




wherein R'_1 and R'_2 may be the same or different and are hydrogen or lower alkyl, $C(O)R_4$ wherein R_4 is independently as defined above, cycloalkyl of from three to twenty carbons having of from three to eight ring carbons, aryl, or aralkyl, (iii) O, or (iv) S; with the proviso that X and Y cannot both be N, NR_5 , O or S at once and with the proviso that one of X and Y cannot be O at the same time the other of X and Y is S or NR_5 and that one of X and Y cannot be S at the same time the other of X and Y is NR_5 ;

(5) Z is H, lower alkyl, aryl, aralkyl, $OC(O)R_3$ wherein R_3 is independently as defined above, $C(O)OR_4$ wherein R_4 is independently as defined above, $C(O)R_3$ wherein R_3 is independently as defined above, $CH(R'_1)CO_2R'_2$ wherein R'_1 and R'_2 are independently as defined above, halogen, trifluoromethyl,



wherein R, R_1 and R_2 are independently as defined above, heteroaryl, or heteroaralkyl; with the overall proviso that when one of R, R_1 , and R_2 is 2-hydroxy, X is O, Y is N and Q is $CH=CH$, then Z cannot be H or alkyl; and also with the overall proviso that when R, R_1 , and R_2 are hydroxy or lower alkyl, Y and X are N or NH, and n is zero then Z cannot be furyl or phenyl unsubstituted or substituted with halogen, trifluoromethyl, alkyl, alkoxy or NO_2 .

The present invention is also a pharmaceutical composition for treating a disease such as allergy, asthma, arthritis, psoriasis, acne, inflammation, pain, or cardiovascular disorders comprising an antiallergic, antiinflammatory, analgesic, or beneficial cardiovascular effective amount of the compound of formula (I)



and pharmaceutically acceptable salts thereof;
wherein

(1) $-$ is a single or double bond;


(2) R, R_1 and R_2 may be the same or different and are selected from the group consisting of hydrogen, lower alkyl, hydroxy, OR_3 wherein R_3 is lower alkyl, $C(O)OR_4$ wherein R_4 is hydrogen or lower alkyl, $OC(O)R_3$ wherein R_3 is independently as defined above, $C(O)R_3$ wherein R_3 is independently as defined above, NR_6R_7 wherein R_6 and R_7 may be the same or different and are hydrogen or lower alkyl, $NHC(O)R_3$ wherein R_3 is independently as defined above, $NHCHO$, $NHSO_2R_3$ wherein R_3 is independently as defined above, $NHCONHR_4$ wherein R_4 is independently as defined above, hydroxymethyl, halogen, trifluoromethyl, SR_4 wherein R_4 is independently as defined above, or nitro;

(3) Q is $(CH_2)_n$ wherein n is an integer of zero to four, $CH=CH$ or




wherein R_4 is independently as defined above;

(4) X and Y are (i) N, (ii) NR₅ wherein R₅ is hydrogen, lower alkyl,



wherein R'₁ and R'₂ may be the same or different and are hydrogen or lower alkyl, C(O)R₃ wherein R₃ is independently as defined above, cycloalkyl of from three to twenty carbons having of from three to eight ring carbons, aryl, or aralkyl, (iii) O, (iv) S; with the proviso that X and Y cannot both be N, NR₅, O or S at once and with the proviso that one of X and Y cannot be O at the same time the other of X and Y is S or NR₅ and that one of X and Y cannot be S at the same time the other of X and Y is NR₅;

(5) Z is H, lower alkyl, aryl, aralkyl, OC(O)R₃ wherein R₃ is independently as defined above, C(O)OR₄ wherein R₄ is independently as defined above, C(O)R₃ wherein R₃ is independently as defined above, CH-(R'₁)CO₂R'₂ wherein R'₁ and R'₂ are independently as defined above, halogen, trifluoromethyl,



wherein R, R₁, and R₂ are independently as defined above, heteroaryl, or heteroaralkyl; with the overall proviso that when R, R₁, and R₂ are hydroxy or lower alkyl, n is zero, Y and X are N or NH, then Z cannot be furyl or phenyl unsubstituted or substituted with halogen, trifluoromethyl, alkyl, alkoxy or NO₂ and a pharmaceutically acceptable carrier.

The present invention is also a composition of the compound of formula I not including the overall provisos and a carrier from among the carriers known to be for use in combination with a sunscreen.

Further, the invention is a method of use of a compound of general formula I or the manufacturing of a pharmaceutical composition for the treatment of diseases noted above.

Also, the invention is a method of using a compound of formula I; again as defined above but not including the overall provisos, as a sunscreen, for example, in a coating on humans, in paint and the like.

Finally, the present invention is a process of preparing a compound of formula I as defined above.

In the compounds of formula I the term lower alkyl is of one to four carbons, inclusive, and includes methyl, ethyl, propyl, or butyl and isomers thereof.

Halogen includes particularly fluorine, chlorine bromine or iodine.

Aryl is phenyl unsubstituted or substituted by one, two or three substituents of one or more of each of alkyl of one to four carbons, inclusive, OR₄ wherein R₄ is independently as defined above, SR₄


wherein R₄ is independently as defined above, R₄ C(=O)O wherein R₄ is independently as defined above, C(O)OR₄ wherein R₄ is independently as defined above, hydroxymethyl, NR₆R₇ wherein R₆ and R₇ are each independently as defined above, or nitro, or halogen.

Aralkyl is an aryl as defined above and attached through an alkylene such as methylene, ethylene, propylene, butylene and isomers thereof.

Heteroaryl means 2-, or 3-pyrrolyl; 2-or 3-furyl; 2-or 3-thiophenyl; 2-, 4-, or 5-oxazoly; 2-, 4-, or 5-thiazoly; 1-, 2-, or 4-imidazoly, 2-, 3-, or 4-isothiazoly, 2-, 3-, or 4-isoxazoly, 1-, 2-, or 3-pyrazoly, and 2-, 3-, 4-pyridyl.

The compounds of the present invention contemplate compounds having the following ring systems

A.




5

10


15

B.




20

25

and
C.



30

35




40

45



50

wherein --Z-- , R_5 and Z are as defined above.When R_5 is hydrogen it is understood the ring system may be represented by the following equilibrium:

55

The compounds of formula I are useful both in the free base form, in the form of base salts where possible, and in the form of acid addition salts. The three forms are within the scope of the invention. In practice, use of the salt form amounts to use of the base form. Appropriate pharmaceutically acceptable salts within the scope of the invention are those derived from mineral acids such as hydrochloric acid and sulfuric acid; and organic acids such as methanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, and the like, giving the hydrochloride, sulfamate, methanesulfonate, benzenesulfonate, p-toluenesulfonate, and the like, respectively or those derived from bases such as suitable organic and inorganic bases. Examples of suitable inorganic bases for the formation of salts of compounds of this invention include the hydroxides, carbonates, and bicarbonates of ammonia, sodium, lithium, potassium, calcium, magnesium, aluminum, zinc, and the like.

Salts may also be formed with suitable organic bases. Bases suitable for the formation of pharmaceutically acceptable base addition salts with compounds of the present invention include organic bases which are nontoxic and strong enough to form such salts. These organic bases form a class whose limits are readily understood by those skilled in the art. Merely for purposes of illustration, the class may be said to include mono-, di-, and trialkylamines, such as methylamine, dimethylamine, and triethylamine; mono-, di- or trihydroxyalkylamines such as mono-, di- and triethanolamine; amino acids such as arginine, and lysine; guanidine; N-methylglucosamine; N-methylglucamine; L-glutamine; N-methylpiperazine; morpholine; ethylenediamine; N-benzylphenethylamine; tris(hydroxymethyl)aminomethane; and the like. (See for example, "Pharmaceutical Salts," J. Pharm. Sci. 66(1):1-19 (1977).)

The acid addition salts of said basic compounds are prepared either by dissolving the free base of compound I, I' or II in aqueous or aqueous alcohol solution or other suitable solvents containing the appropriate acid or base and isolating the salt by evaporating the solution, or by reacting the free base of compound I, I' or II with an acid as well as reacting compound I, I' or II having an acid group thereon with a base such that the reactions are in an organic solvent, in which case the salt separates directly or can be obtained by concentration of the solution.

The compounds of the invention may contain an asymmetric carbon atom. Thus, the invention includes the individual stereoisomers, and the mixtures thereof. The individual isomers may be prepared or isolated by methods known in the art.

Compounds of the present invention that are preferred are of formula I wherein Y is N and X is NH or Y is N and X is O.

The most preferred compounds of the present invention are 5-[β -(4'-hydroxy-3',5'-bis(1,1-dimethylethyl)-phenyl)ethenyl]-3-methylisoxazole, 3-[β -(4'-hydroxy-3'-methoxyphenyl)ethenyl]-5-methylpyrazole, and 5-[β -(4'-hydroxy-3',5'-dimethoxyphenyl)ethenyl]-3-methylisoxazole.

A physician or veterinarian of ordinary skill readily determines a subject who is exhibiting symptoms of any one or more of the diseases described above. Regardless of the route of administration selected, the compounds of the present invention of the formula I as described in pharmaceutical compositions above are formulated into pharmaceutically acceptable dosage forms by conventional methods known to the pharmaceutical art.

The compounds can be administered in such oral unit dosage forms as tablets, capsules, pills, powders, or granules. They also may be administered rectally or vaginally in such forms as suppositories or bougies; they may also be introduced parenterally (e.g. subcutaneously, intravenously, or intramuscularly), using forms known to the pharmaceutical art. They are also introduced directly to an affected area (e.g., in the form of eye drops or by inhalation). For the treatment of asthma or allergies, particularly dermatological disorders; such as erythema, psoriasis and acne, the compounds may also be administered topically in the form of ointments, gels, or the like. However, in general, the preferred route of administration is orally.

An effective but nontoxic quantity of the compound is employed in treatment. The ordinarily skilled physician or veterinarian will readily determine and prescribe the effective amount of the compound to prevent or arrest the progress of the condition for which treatment is administered. In so proceeding, the physician or veterinarian could employ relatively low dosages at first, subsequently increasing the dose until a maximum response is obtained.

Initial dosages of the compounds of formula I of the invention in pharmaceutical compositions are ordinarily in the area of 10 mg up to 2 grams per day orally, preferably 10 mg to 500 mg per dose orally given one to four times daily or as needed. When other forms of administration are employed equivalent doses are administered.

Additionally, the present invention is compositions comprising the compounds of formula I, for use as sunscreens, having a suitable carrier therefor.

The ultraviolet absorbing properties of the compounds of the present invention is generally shown by a comparison with p-aminobenzoic acid which is the active ingredient in most commercial sunscreens. Such properties are within the ultraviolet absorbing ranges of 260 to 300 nM range critical to effectiveness. A representative compound of the invention is compared to p-aminobenzoic acid in the following Table.


TABLE

<u>Compound</u>	<u>UV max</u>	<u>α</u>
p-aminobenzoic acid	285 nM	132.2
3-[β -(4'-hydroxy-3'-methoxy-phenyl)ethenyl]-5-methylpyrazole	292 nM	111.1

The methods of preparation for the compounds of the present invention may, generally, be accomplished in a manner according to the definition of X and Y.

For example, generally compounds of formula I wherein X and Y are each N or NR₅ are prepared in a manner shown by Scheme I as follows:

SCHEME I




wherein --- , R, R₁, R₂, R₅ and Z are as defined above.

In Step 1 of Scheme I equimolar parts of hydrazine hydrate and a compound of the formula III₃ wherein R, R₁, R₂ and Z are as defined above are combined in a solvent such as 1 part ethanol and 1 part butanol, or 2-propanol, and the like. A small amount of about 0.1 ml to 10 ml preferably 0.5 ml of acetic acid is added with the hydrazine hydrate. Optionally, the step 1 product of formula I₁ is further reacted as shown in

step II with a compound MR_5 wherein M is an electrofuge such as halogen, sulfonate, or the like and R_5 is as defined above but other than hydrogen to obtain a mixture of the compound of formula I_1' and I_1'' wherein --- , R, R_1 , R_2 , R_5 when not hydrogen and Z are as defined above. The compounds I_1' and I_1'' may be separated by conventional means.







Similarly compounds of formula I wherein X and Y are each N or NR_5 wherein Q is CH_2 and n is 1 can be prepared according to the following Scheme Ia using analogous conditions as provided above for Scheme I.

SCHEME Ia




Further, the compounds of formula I_2' are prepared as shown in the following Scheme II and IIa.

SCHEME II


 III_2  II_2  I'_2

optionally
 H_2

 I_2

wherein X and Y are O or N and S or N .

SCHEME IIa



Triphenylphosphonium halide; preferably chloride, of the formula II₂ or II_{2a} are contacted by an excess of a compound of the formula III₂ or III_{2a} in a solvent such as dimethylsulfoxide, toluene, or the like to which an equivalent amount of potassium t-butoxide is added. The mixture is stirred at about room temperature for from 10 minutes to 2 hours or more under an inert atmosphere. A compound of formula I₂^u or I_{2a}^u is obtained wherein R, R₁, R₂ and Z are as defined above and Y and X are each either N or O and N or S, respectively.

Mild hydrogenation by conventional methods by an ordinarily skilled artisan may be carried out to provide a compound of formula I₂^u wherein R, R₁, R₂ and Z are as defined and Y and X are O or N and S or N, respectively.

Additionally, for compounds wherein Z is defined such that the two substituents on the heterocyclic ring of formula


45



50


or

55




are the same, the compounds may be prepared according to the following Scheme III.

SCHEME III





wherein Z is



, and wherein R, R₁, R₂, and Q are as defined above and Y and X are either O and N or N and O.

The conditions of the reaction in Scheme III are, generally, carried out with equimolar parts of the compound of formula III₁ and hydroxylamine hydrochloride in a solvent such as methanol, ethanol and the like in the presence of a buffer such as sodium acetate at reflux temperature until completion as determined by TLC.


SCHEME IV

AB

wherein R, R_1 , R_2 , and Z are as defined above.

The conditions of the synthesis in Scheme IV step 1 are, generally, carried out in a manner analogous to those described by C. Kashima et al, Bull. Chem. Soc. Japan, **46**, 310 (1973) and C. Kashima et al, Heterocycles, **6**, 805 (1977). Unexpectedly the step 2 dehydration when R₄ is H or alkyl and also further decarboxylation when R₄ is H are carried out in a one pot reaction together with step 1. Conditions for step 2 are analogous to those known in the art and, thus, within the ordinary artisan's skill.


SCHEME V




The conditions of the above Scheme V are analogous to those described by M. Nitta, J. Chem. Soc. Chem. Commun., 877, 1982.

Finally, conditions analogous to those described in copending application PD-3493 United States Serial Number 851,003, filed April 11, 1986 are useful in step 1 of Scheme VI below. Step 2 of Scheme VI is carried out using anhydrous conditions in an inert solvent at about the temperature of 0° to 50°C for from 90 min to 18 hours.

SCHEME VI




SCHEME VII



The compounds in Scheme VII are compounds wherein R, R₁, R₂ and Z are as defined above for compounds of formula III₃' and III₃' but III₁' and III₁' are symmetrical.


An alternate preparation for the compounds of formula III₃ is as found in Scheme VIII as follows:

SCHEME VIII



Finally, the compound of formula II_2 as shown in Scheme II above can be prepared by the following method.

SCHEME IX



Under certain circumstances it is necessary to protect either the N or O of intermediates in the above noted process with suitable protecting groups which are known. Introduction and removal of such suitable oxygen and nitrogen protecting groups are well-known in the art of organic chemistry; see for example "Protective Groups in Organic Chemistry," J. F. W. McOmie, ed., (New York, 1973), pages 43ff, 95ff, J. F. W. McOmie, *Advances in Organic Chemistry*, Vol. 3, 191-281 (1963); R. A. Borssenas, *Advances in Organic Chemistry*, Vol. 3, 159-190 (1963); and J. F. W. McOmie, *Chem. & Ind.*, 603 (1979).

Examples of suitable oxygen protecting groups are benzyl, t-butyldimethylsilyl, ethoxyethyl, and the like. Protection of an N-H containing moiety is necessary for some of the processes described herein for the preparation of compounds of this invention. Suitable nitrogen protecting groups are benzyl, triphenylmethyl, trialkylsilyl, trichloroethylcarbamate, trichloroethoxycarbonyl, vinylloxycarbamate, and the like.

Under certain circumstances it is necessary to protect two different oxygens with dissimilar protecting groups such that one can be selectively removed while leaving the other in place. The benzyl and t-butyldimethylsilyl groups are used in this way; either is removable in the presence of the other, benzyl being removed by catalytic hydrogenolysis, and t-butyldimethylsilyl being removed by reaction with, for example, tetra-n-butylammonium fluoride.

In the process described herein for the preparation of compounds of this invention the requirements for protective groups are generally well recognized by one skilled in the art of organic chemistry, and accordingly the use of appropriate protecting groups is necessarily implied by the processes of the charts herein, although not expressly illustrated.

The products of the reactions described herein are isolated by conventional means such as extraction, distillation, chromatography, and the like.

Starting materials not described herein are available commercially, are known, or can be prepared by methods known in the art.

The salts of the compounds of formula I described above are prepared by reacting the appropriate base or acid with a stoichiometric equivalent of the compounds of formula I, respectively, to obtain pharmaceutically acceptable salts thereof.

The invention is further elaborated by the representative examples as follows. Such examples are not meant to be limiting.

Example 13,5-Bis[β -(4'-hydroxy-3'-methoxyphenyl)ethenyl]pyrazole

5 Curcumin (5.93 g, 16.1 mmoles) (Sigma Co., St. Louis) is dissolved in 100 ml of 50% ethanol in butanol. Hydrazine hydrate (0.81 g, 16.1 mmoles) and 0.5 ml of acetic acid are added. The reaction is warmed in an oil bath at 60°C for 24 hours. The solvents are removed under vacuum, and the residue is chromatographed on silica gel in ethyl acetate to give a red solid. Recrystallization of the product from methanol/water gives 0.3 g of the product as a hydrate. Analysis for $C_{21}H_{20}N_2O_4 \cdot 0.1$ mole H_2O requires C-68.87, H-5.52, N-7.65. Found; C-68.77, H-5.50, N-7.44. mp = 211-214°C.

Example 215 3,5-Bis[β -(4'-hydroxy-3'-methoxyphenyl)-5-ethylpyrazole

According to the procedure of Example 1, 1,7-bis(4'-hydroxy-3'-methoxyphenyl)-3,5-heptadione (Bull. Acad. Polon. Sci. 6, 481-486, 1958) is reacted with hydrazine hydrate to afford 3,5-Bis[β -(4'-hydroxy-3'-methoxyphenyl)ethyl]pyrazole in 65% yield, mp = 125-128°C.

Example 325 3,5-Bis[β -(4'-hydroxyphenyl)ethyl]pyrazole

According to the procedure of Example 1, 1,7-bis(4'-hydroxyphenyl)-3,5-heptadione is reacted with hydrazine hydrate to afford 3,5-bis[β -(4'-hydroxyphenyl)ethyl]pyrazole in 45% yield, mp = 193-195°C.

30 Example 43-[β -(4'-Hydroxy-3'-methoxyphenyl)ethenyl]-5-phenylpyrazole

According to the procedure of Example 1, 1-(4'-hydroxy-3'-methoxyphenyl)-5-phenyl-1-pentene-3,5-dione is reacted with hydrazine hydrate to afford 3-[β -(4'-hydroxy-3'-methoxyphenyl)ethenyl]-5-phenylpyrazole in 9% yield, mp = 132-134°C.

Example 540 3-[β -(4'-Hydroxy-3'-methoxyphenyl)ethenyl]-5-methylpyrazole

According to the procedure of Example 1, 1-(4'-hydroxy-3'-methoxyphenyl)-1-hexene-3,5-dione is reacted with hydrazine hydrate to afford 3-[β -(4'-hydroxy-3'-methoxyphenyl)ethenyl]-5-methylpyrazole in 53% yield, mp = 144-146°C.

Example 650 3-[β -(4'-hydroxy-3'-methoxyphenyl)ethyl]-5-methylpyrazole

According to the procedure of Example 1, 1-(4'-hydroxy-3'-methoxyphenyl)-3,5-hexanedione is reacted with hydrazine hydrate to afford 3-[β -(4'-hydroxy-3'-methoxyphenyl)ethyl]-5-methylpyrazole in 68% yield, mp = 84-86°C.

Example 7

3-[β -(4'-Hydroxy-3'-chlorophenyl)ethenyl]-5-methyl-pyrazole

According to the procedure of Example 1, 1-(4'-hydroxy-3'-chlorophenyl)-1-hexene-3,5-dione is reacted with hydrazine hydrate to afford 3-[β -(4'-hydroxy-3'-chlorophenyl)ethenyl]-5-methyl-pyrazole in 61% yield, mp = 183-184°C.

Example 83-[β -(4'-Methoxy-3'-hydroxyphenyl)ethenyl]-5-methyl-pyrazole

According to the procedure of Example 1, 1-(4'-methoxy-3'-hydroxyphenyl)-1-hexene-3,5-dione is reacted with hydrazine hydrate to afford 3-[β -(4'-methoxy-3'-hydroxyphenyl)ethenyl]-5-methyl-pyrazole in 72% yield, mp = 221-222°C.

Example 93-[β -(4'-Hydroxy-3'-methoxyphenyl)ethenyl]-5-ethyl-pyrazole

According to the procedure of Example 1, 1-(4'-hydroxy-3'-methoxyphenyl)-1-heptene-3,5-dione is reacted with hydrazine hydrate to afford 3-[β -(4'-hydroxy-3'-methoxyphenyl)ethenyl]-5-ethyl-pyrazole in 26% yield, mp = 61-64°C.

Example 10Ethyl-3-[β -(3-methoxy-4-hydroxyphenyl)ethenyl]-1H-pyrazole-5-carboxylate

According to the procedure of Example 1, ethyl 6-(3-methoxy-4-hydroxyphenyl)-2,4-dioxo-5-hexenoate is reacted with hydrazine hydrate to afford ethyl-3-[β -(3-methoxy-4-hydroxyphenyl)ethenyl]-1H-pyrazole-5-carboxylate in 51% yield, mp = 99-101°C.

Example 10A3-[β -(4'-Hydroxy-3'-methoxyphenyl)ethenyl]-5-trifluoromethylpyrazole

According to the procedure of Example 1, 1-(4'-hydroxy-3'-methoxyphenyl)-6,6,6-trifluoro-1-hexene-3,5-dione is reacted with hydrazine hydrate to afford 3-[β -(4'-hydroxy-3'-methoxyphenyl)ethenyl]-5-trifluoromethylpyrazole in 71% yield, mp = 145-147°C.

Example 10B3-[β -(4'-Butyldimethylsilyloxy-3'-methoxyphenyl)ethenyl]-5-methylpyrazole

According to the procedure of Example 1, 1-(4'-t-butyldimethylsilyloxy-3'-methoxyphenyl)-1-hexene-3,5-dione is reacted with hydrazine hydrate to afford 3-[β -(4'-t-butyldimethylsilyloxy-3'-methoxyphenyl)ethenyl]-5-methylpyrazole in 52% yield, sufficiently pure for further use.

Example 10C

3-[β -(4'-Butyldimethylsilyloxy-3,5-dimethoxyphenyl)ethenyl]-5-methylpyrazole

According to the procedure of Example 1, 1-(4'-t-butyldimethylsilyloxy-3',5'-dimethoxyphenyl)-1-hexene-3,5-dione is reacted with hydrazine hydrate to afford 3-[β -(4'-t-butyldimethylsilyloxy-3',5'-dimethoxyphenyl)-ethenyl]-5-methylpyrazole in 32% yield, mp = 133-136°C.

Example 10D10 (E)-5-[β -(4'-Hydroxy-3',5'-dimethoxyphenyl)ethenyl]-3-methyl-1H-pyrazole-1-acetic acid, ethyl ester

A solution of 1-(4'-hydroxy-3',5'-dimethoxyphenyl)-1-hexene-3,5-dione (1.0 g, 3.8 mmol) in absolute ethanol (100 mL) is added to a solution of ethyl hydrazinoacetate in absolute ethanol (100 mL) according to the procedure found in A. Carmi et al, *J. Org. Chem.*, 25, 44 (1960). The reaction mixture is acidified with acetic acid (2 mL) and is stirred for 30 minutes at room temperature. The solvent is evaporated, and the residue is suspended in water (50 mL). The solid which is collected is then purified by flash chromatography (silica gel, 1:1 methylenechloride/ethyl acetate) and subsequent recrystallization (methylenechloride/ethyl acetate 1:1) to afford 0.45 g (34%) of (E)-5-[β -(4'-hydroxy-3',5'-dimethoxyphenyl)-ethenyl]-3-methyl-1H-pyrazole-1-acetic acid, ethyl ester, mp = 159-160°C. Analysis for C₁₈H₂₂N₂O₅. Calculated C-62.42, H-6.40, N-8.09. Found C-62.08, H-6.39, N-8.26.

Example 10E25 5-[β -(4'-t-Butyldimethylsilyloxy-3'-methoxyphenyl)ethenyl]-3-methyl-1H-pyrazole-1-acetic acid, methyl ester; and 3-[β -(4'-t-butyldimethylsilyloxy-3'-methoxyphenyl)ethenyl]-5-methyl-1H-pyrazole-1-acetic acid, methyl ester

3-[β -(4'-t-butyldimethylsilyloxy-3'-methoxyphenyl)ethenyl]-5-methylpyrazole (2.57 g, 7.5 mmol) is dissolved in DMF (20 mL). Potassium carbonate (10.30 g, 75 mmol) is ground into a fine powder and added to the reaction. The reaction is stirred at room temperature for 10 minutes under argon atmosphere. Methyl bromoacetate is added and the reaction is stirred for 2 hours. The mixture is then poured into water (200 mL) and neutralized with aqueous HCl. The layers are separated, and the aqueous layer is washed with methylenechloride. The combined organic layers are washed with water and brine. Drying over MgSO₄ followed by evaporation of solvents affords a brown oil. Flash chromatography (10% ethyl ether in methylenechloride) separates two products: 3-[β -(4'-t-butyldimethylsilyloxy-3'-methoxyphenyl)ethenyl]-5-methyl-1H-pyrazole-1-acetic acid, methyl ester (R_f = 0.31); 1.5 g; and 5-[β -(4'-t-butyldimethylsilyloxy-3'-methoxyphenyl)ethenyl]-3-methyl-1H-pyrazole-1-acetic acid, methyl ester (R_f = 0.14), 0.55 g.

Example 10F45 3-[β -(4'-t-butyldimethylsilyloxy-3',5'-dimethoxy phenyl)ethenyl]-5-methyl-1H-pyrazole-1-acetic acid, methyl ester and 5-[β -(4'-t-butyldimethylsilyloxy-3',5'-dimethoxyphenyl)ethenyl]-3-methyl-1H-pyrazole-1-acetic acid, methyl ester

According to the procedure of Example 10E, 3-[β -(4'-t-butyldimethylsilyloxy-3',5'-dimethoxyphenyl)-ethenyl]-5-methylpyrazole (3.5 g, 9.35 mmol) is reacted with methyl bromoacetate to afford 3-[β -(4'-t-butyldimethylsilyloxy-3',5'-dimethoxyphenyl)ethenyl]-5-methyl-1H-pyrazole-1-acetic acid, methyl ester, 2.3 g (55%), mp = 139-141°C; and 5-[β -(4'-t-butyldimethylsilyloxy-3',5'-dimethoxyphenyl)ethenyl]-3-methyl-1H-pyrazole-1-acetic acid, methyl ester, 0.9 g (21%), mp = 150-151°C.

Example 10G

3-[β -(4'-Hydroxy-3',5'-di-t-butylphenyl)ethenyl]-5-methyl-1H-pyrazole-1-acetic acid, methyl ester and 5-[β -(4'-hydroxy-3',5'-di-t-butylphenyl)ethenyl]-3-methyl-1H-pyrazole-1-acetic acid, methyl ester

According to the procedure of Example 10E, 3-[β -(4'-hydroxy-3',5'-di-t-butylphenyl)ethenyl]-5-methyl-
 5 pyrazole (3.0 g, 9.6 mmol) is reacted with methyl bromoacetate in DMF, with the exception that sodium acetate is used in place of potassium carbonate. Flash chromatography separates 3-[β -(4'-hydroxy-3',5'-di-t-butylphenyl)ethenyl]-5-methyl-1H-pyrazole-1-acetic acid, methyl ester, 0.65 g (18%), mp = 170-174°C; and 5-[β -(4'-hydroxy-3',5'-di-t-butylphenyl)ethenyl]-3-methyl-1H-pyrazole-1-acetic acid, methyl ester, (as HCl salt) 0.72 g (17%), mp = 179-180°C. Analysis for $C_{24}H_{33}ClN_2O_3$. Calculated C-65.62, H-7.90, N-6.65, Cl-8.42.
 10 Found C-65.53, H-7.99, N-6.46, Cl-8.31.

Example 10G'

15 3-[β -(4'-Hydroxy-3',5'-di-t-butylphenyl)ethenyl]-1,5-dimethylpyrazole and 5-[β -(4'-hydroxy-3',5'-di-t-butylphenyl)ethenyl]-1,3-dimethylpyrazole

According to the procedure of Example 10G, 3-[β -(4'-hydroxy-3',5'-di-t-butylphenyl)ethenyl]-5-methyl-
 20 pyrazole is reacted with methyl iodide and sodium acetate to afford two regioisomeric products separated by flash chromatography (chloroform): 3-[β -(4'-hydroxy-3',5'-di-t-butylphenyl)ethenyl]-1,5-dimethylpyrazole, mp = 155-157°C; 5-[β -(4'-hydroxy-3',5'-di-t-butylphenyl)ethenyl]-1,3-dimethylpyrazole, mp = 120-122°C.

Example 10H

25 3-[β -(4'-Hydroxy-3',5'-dimethoxyphenyl)ethenyl]-5-methyl-1H-pyrazole-1-acetic acid, methyl ester

To a solution of 3-[β -(4'-t-butylidimethylsilyloxy-3',5'-dimethoxyphenyl)ethenyl]-5-methyl-1H-pyrazole-1-
 30 acetic acid, methyl ester (2.25 g, 5.0 mmol) in dry THF (150 mL) is added tetra-n-butyl-ammonium fluoride (7.5 mL, 1M solution in THF) at 0°C. The reaction mixture is stirred at 0°C for one hour and then quenched by the dropwise addition of brine (100 mL) at 0°C. The layers are separated and the aqueous layer extracted with ethyl acetate. The combined organic layers are evaporated and the residue purified by flash chromatography (ethyl acetate/methylene chloride 1:1) to afford 1.35 g (80%) of 3-[β -(4'-hydroxy-3',5'-dimethoxyphenyl)ethenyl]-5-methyl-1H-pyrazole-1-acetic acid methyl ester, mp = 112-114°C. Analysis for $C_{17}H_{20}N_2O_5$. Calculated C-61.44, H-6.07, N-8.43. Found C-61.23, H-6.08, N-8.44.
 35

Example 10I

40 5-[β -(4'-Hydroxy-3'-methoxyphenyl)ethenyl]-3-methyl-1H-pyrazole-1-acetic acid methyl ester

According to the procedure of Example 10H, 5-[β -(4'-t-butylidimethylsilyloxy-3'-methoxyphenyl)ethenyl]-
 3-methyl-1H-pyrazole-1-acetic acid methyl ester is reacted with tetra-n-butyl ammonium fluoride to afford 5-
 45 [β -(4'-hydroxy-3'-methoxyphenyl)ethenyl]-3-methyl-1H-pyrazole-1-acetic acid methyl ester, in 46% yield, mp = 159-160°C. Analysis for $C_{16}H_{18}N_2O_4$. Calculated C-63.55, H-6.01, N-9.27. Found C-63.68, H-6.05, N-9.20.

Example 10J

50 3-[β -(4'-Hydroxy-3'-methoxyphenyl)ethenyl]-5-methyl-1H-pyrazole-1-acetic acid, methyl ester

According to the procedure of Example 10H, 3-[β -(4'-t-butylidimethylsilyloxy-3'-methoxyphenyl)ethenyl]-
 5-methyl-1H-pyrazole-1-acetic acid methyl ester is reacted with tetra-n-butyl ammonium fluoride to afford 3-
 55 [β -(4'-hydroxy-3'-methoxyphenyl)ethenyl]-5-methyl-1H-pyrazole-1-acetic acid methyl ester, in 78% yield, mp = 141-143°C. Analysis for $C_{16}H_{18}N_2O_4$. Calculated C-63.55, H-6.01, N-9.27. Found C-63.53, H-6.02, N-8.89.

Example 10K

3-[β -(4'-Hydroxy-3'-methoxyphenyl)ethenyl]-5-methyl-1H-pyrazole-1-acetic acid

A solution of 3-[β -(4'-hydroxy-3'-methoxyphenyl) ethenyl]-5-methyl-1H-pyrazole-1-acetic acid methyl ester (1.1 g, 3.6 mmol) in aqueous sodium hydroxide (200 mL, 1N solution) with the minimum amount of hot ethanol needed for dissolution, is heated on a steam bath for 2 hours. The reaction mixture is cooled and neutralized with 4 N HCl. The resulting precipitate is collected by filtration and recrystallized from 2-propanol/water to afford 3-[β -(4'-hydroxy-3'-methoxyphenyl)ethenyl]-5-methyl-1H-pyrazole-1-acetic acid, 0.64 g (62%), mp = 225-232°C. Analysis for $C_{15}H_{16}N_2O_4$. Calculated C-62.49, H-5.59, N-9.72. Found C-62.35, H-5.64, N-9.69.

Example 10L5-[β -(4'-Hydroxy-3'-methoxyphenyl)ethenyl]-3-methyl-1H-pyrazole-1-acetic acid

According to the procedure of Example 10K, 5-[β -(4'-hydroxy-3'-methoxyphenyl)ethenyl]-3-methyl-1H-pyrazole-1-acetic acid methyl ester is saponified to afford 5-[β -(4'-hydroxy-3'-methoxyphenyl)ethenyl]-3-methyl-1H-pyrazole-1-acetic acid, in 71% yield. Mp = 265-268°C. Analysis for $C_{15}H_{16}N_2O_4$. Calculated C-62.49, H-5.59, N-9.72. Found C-62.26, H-5.58, N-9.56.

Example 10M5-[β -(4'-Hydroxy-3',5'-dimethoxyphenyl)ethenyl]-3-methyl-1H-pyrazole-1-acetic acid

According to the procedure of Example 10K, 5-[β -(4'-Hydroxy-3',5'-dimethoxyphenyl)ethenyl]-3-methyl-1H-pyrazole-1-acetic acid methyl ester is saponified to afford 5-[β -(4'-hydroxy-3',5'-dimethoxyphenyl)ethenyl]-3-methyl-1H-pyrazole-1-acetic acid, in 90% yield. Mp = 240-243°C. Analysis for $C_{16}H_{18}N_2O_5 \cdot 0.1 H_2O$. Calculated C-60.03, H-5.73, N-8.75. Found C-59.90, H-5.50, N-8.66.

Example 10N3-[β -(4'-Hydroxy-3',5'-dimethoxyphenyl)ethenyl]-5-methyl-1H-pyrazole-1-acetic acid

According to the procedure of Example 10K, 3-[β -(4'-Hydroxy-3',5'-dimethoxyphenyl)ethenyl]-5-methyl-1H-pyrazole-1-acetic acid methyl ester is saponified to afford 3-[β -(4'-hydroxy-3',5'-dimethoxyphenyl)ethenyl]-5-methyl-1H-pyrazole-1-acetic acid, in 65% yield. Mp = 232-234°C. Analysis for $C_{16}H_{18}N_2O_5$. Calculated C-60.37, H-5.70, N-8.80. Found C-60.20, H-5.88, N-8.53.

Example 10P5-[β -(4'-Hydroxy-3',5'-di-t-butylphenyl)ethenyl]-3-methyl-1H-pyrazole-1-acetic acid

According to the procedure of Example 10K, 5-[β -(4'-hydroxy-3',5'-di-t-butylphenyl)ethenyl]-3-methyl-1H-pyrazole-1-acetic acid methyl ester is saponified to afford 5-[β -(4'-hydroxy-3',5'-di-t-butylphenyl)ethenyl]-3-methyl-1H-pyrazole-1-acetic acid, in 50% yield. Mp = 220-224°C. Analysis for $C_{22}H_{30}N_2O_3$. Calculated C-71.32, H-8.16, N-7.56. Found C-71.14, H-8.16, N-7.46.

Example 10Q

3-[β -(4'-Hydroxy-3',5'-di-*t*-butylphenyl)ethenyl]-5-methyl-1H-pyrazole-1-acetic acid

According to the procedure of Example 10K, 3-[β -(4'-hydroxy-3',5'-di-*t*-butylphenyl)ethenyl]-5-methyl-1H-pyrazole-1-acetic acid methyl ester is saponified to afford 3-[β -(4'-hydroxy-3',5'-di-*t*-butylphenyl)ethenyl]-5-methyl-1H-pyrazole-1-acetic acid, in 50% yield. Mp = 227-229°C. Analysis for $C_{22}H_{30}N_2O_3$. Calculated C-71.32, H-8.16, N-7.56. Found C-71.22, H-8.18, N-7.53.

Example 11

3-[β -(3-Methoxy-4-hydroxyphenyl)ethenyl]-1H-pyrazole-5-carboxylic acid

Ethyl-3-[β -(3-methoxy-4-hydroxyphenyl)ethenyl]-1H-pyrazole-5-carboxylate (0.49 g, 1.7 mmoles) is added to a solution of KOH (0.38 g, 6.8 mmoles) in 50 ml EtOH. The reaction is warmed to reflux overnight. After cooling to room temperature, the reaction is diluted to 200 ml with H_2O , and acidified to pH=4 with HCl. A white precipitate forms, and it is collected by filtration. Drying in vacuum in the presence of P_2O_5 gives 0.3 g (68%) of the desired product, mp = 270-271°C (dec).

Example 12

3,5-Bis[β -(4'-hydroxy-3'-methoxyphenyl)ethyl]-isoxazole

1,7-Bis(4'-hydroxy-3'-methoxyphenyl)-3,5-heptadione (1.00 g, 2.7 mmoles) (Bull. Acad. Polon. Sci. 6, 481-486, 1958), hydroxylaminehydrochloride (0.19 g, 2.7 mmoles), and sodium acetate (0.44 g, 5.4 mmoles) are dissolved in 50 ml of 95% ethanol. The reaction is stirred at reflux overnight. The solvent is removed, and the residue is taken up in ethyl acetate. The ethyl acetate solution is washed with water and dried over magnesium sulfate. A precipitate forms on standing at room temperature for 2 days, and it is collected by filtration to give a white solid. The solid is taken up in 20 ml of ethanol containing 0.5 ml of 2N hydrochloric acid. The solution is heated to reflux for one hour. After neutralization with sodium bicarbonate, the ethanol is evaporated, and the residue is partitioned between ethyl acetate and water. The organic layer is dried over magnesium sulfate and passed through a silica gel plug which is washed with 1:1 ethyl acetate/hexane. Evaporation of the solvent gives 0.2 g of the product obtained as a hydrate. mp = 106-107°C. Analysis for $C_{21}H_{25}NO_6 \cdot 0.4$ mole H_2O . Calculated C-66.97, H-6.38, N-3.72. Found C-67.13, H-6.41, N-3.47.

Example 13

5-[β -(4'-Acetoxy-3'-methoxyphenyl)ethenyl]-3-trifluoromethylisoxazole

To a solution of isoxazole-3-trifluoromethyl-5-methyltriphenylphosphonium chloride (350 mg, 0.78 mmol) in dimethylsulfoxide (5 ml) is added potassium *t*-butoxide (120 mg, 1.10 mmol). The mixture is stirred at room temperature for 45 minutes under an inert atmosphere. Acetylvanillin (170 mg, 0.89 mmol) is then added and the reaction stirred an additional one hour. The mixture is poured into saturated aqueous ammonium chloride and extracted into ethyl acetate. The organic layer is washed with water and dried over magnesium sulfate. Concentration and flash chromatography (hexane/ethyl acetate 3:1) afforded 150 mg of 5-[β -(4'-acetoxy-3'-methoxyphenyl)ethenyl]-3-trifluoromethylisoxazole, of sufficient purity for subsequent reaction.

Example 13A

(E)-5-[β -(4'-Hydroxy-3',5'-bis(1,1-dimethylethyl) phenyl)ethenyl]-3-trifluoromethylisoxazole

According to the procedure of Example 13, potassium *t*-butoxide (264 mg), isoxazole-3-trifluoromethyl-5-methyl triphenylphosphonium chloride (1.054 g), and 3,5-di-*t*-butyl-4-hydroxybenzaldehyde (250 mg) are reacted in DMSO for 24 hr at room temperature to afford a *cis*-/*trans*-mixture of two olefinic products. Refluxing a mixture of this isomeric mixture in HCl/methanol (12 hrs) affords pure (E)-5-[β -(4'-hydroxy-3',5'-bis(1,1-dimethylethyl)phenyl)ethenyl]-3-trifluoromethylisoxazole (350 mg); mp = 132-134°C. Analysis for $C_{20}H_{24}NO_2F_3$. Calculated C-65.38, H-6.58, N-3.81. Found C-65.58, H-6.68, N-3.92.

Example 145-[β -(4'-Hydroxy-3'-methoxyphenyl)ethenyl]-3-trifluoromethylisoxazole

A solution of 5-[β -(4'-acetoxy-3'-methoxyphenyl)ethenyl]-3-trifluoromethylisoxazole (120 mg, 0.34 mmol) in methanol (3 ml) is cooled to 0°C. Sodium methoxide (40 mg, 0.68 mmol) is added and the mixture stirred for 30 minutes. The mixture is diluted with ethyl acetate and washed with water. The organic layer is dried (magnesium sulfate) and concentrated. Recrystallization from hexane/ethyl acetate affords 80 mg of pure 5-[β -(4'-hydroxy-3'-methoxyphenyl)ethenyl]-3-trifluoromethylisoxazole; yield 82%; mp = 83-87°C.

Example 151,7-Bis-(4'-hydroxyphenyl)-3,5-heptadione

A solution of 1,7-bis-(4'-hydroxyphenyl)-1,6-heptadiene-3,5-dione (1.99 g, 6.5 mmoles) in tetrahydrofuran (100 ml) is hydrogenated over Raney-nickel (0.2 g) under 30.4 psi of H_2 . The THF is evaporated, and the residue chromatographed in 5% MeOH/ $CHCl_3$ to give 1.6 g (79%) of the desired product, mp = 108-110°C.


Example 161-(3'-Methoxy-4'-hydroxyphenyl)hexene-3,5-dione


According to the procedure of Example 15, 1-(3'-methoxy-4'-hydroxyphenyl)-1-hexene-3,5-dione is hydrogenated to afford in 90% yield 1-(3'-methoxy-4'-hydroxyphenyl)hexene-3,5-dione as an oil.


Example 171-(4'-Hydroxy-3'-methoxyphenyl)-1-hexene-3,5-dione

Boric anhydride (16.32 g, 237.2 mmoles) and acetylacetone are stirred at room temperature under nitrogen for 12 hours. 4-Hydroxy-3-methoxybenzaldehyde (18.06 g, 118.8 mmoles) and tributyl borate (364 g, 3.16 moles) are combined with the acetyl acetone/boric anhydride complex in 100 ml of ethyl acetate. The reaction is stirred at room temperature for one hour, at which time *n*-butylamine (4.34 g, 59.4 mmoles) is added in 4 equal portions over 8 hours. The reaction is allowed to stir for 12 hours. 300 ml of water containing 20 ml of concentrated hydrochloric acid is added to the reaction and stirred for 2 hours. The solid which forms is removed by filtration through celite, and the layers of filtrate are separated. The ethyl acetate layer is washed with 2 \times 100 ml of saturated sodium bicarbonate, followed by 100 ml of brine. Drying over magnesium sulfate followed by evaporation of the solvent gives a red solid. Chromatography in 5% ethyl acetate/chloroform gives 11.06 g (40%) of a yellow solid. mp = 146-147°C. Analysis for $C_{18}H_{14}O_4$ requires C-66.65, H-6.04. Found C-66.88, H-6.18.

The following compounds were made by the above procedure:



Ex.	R	R ₁	Z	Melting Point	Yield
18	HO	Cl	Me	155-157°C	14%
19	MeO	HO	Me	160-162°C	13%
20	HO	MeO	 ¹	155-157°C	75%
21	HO	MeO	Et ²	101-102°C	21%

Example 22Ethyl 6-(3'-methoxy-4'-hydroxyphenyl)-2,4-dioxo-5-hexenoate

Sodium metal (1.8 g, 0.078 mole) is dissolved in absolute ethanol (50 ml) under an inert atmosphere. An ethanolic solution of 1-(3'-methoxy-4'-hydroxyphenyl)-1-butene-3-one (5.00 g, 0.026 mole) is then added slowly to the sodium ethoxide solution. The mixture is stirred for 10 minutes, after which diethyl oxalate (3.80 g, 0.026 mole) is added. The reaction is stirred at room temperature for 5 hours and then acidified with concentrated HCl. The mixture is then stirred at 0°C for 1 hour. The precipitate is collected and dried to afford 4.8 g of ethyl 6-(3'-methoxy-4'-hydroxyphenyl)-2,4-dioxo-5-hexenoate; 63% yield; mp = 97-98°C.

Example 22A1-(4'-t-Butyldimethylsilyloxy-3'-methoxyphenyl)-1-hexene-3,5-dione

A solution of 1-(4'-t-butyldimethylsilyloxy-3'-methoxyphenyl)-1-butene-3-one (12.8 g, 41.8 mmol) in THF (50 mL) is cooled to -78°C under an argon atmosphere. Lithium di-isopropyl amide (83.5 mmol, in 50 mL THF) is slowly added to the reaction mixture. The reaction is stirred at -78°C for 30 minutes. Acetyl chloride (16.4 g, 209 mmol) dissolved in THF (20 mL) is then added to the reaction over a 30 minute period. The

reaction is quenched with aqueous HCl. The layers are separated, and the aqueous layer is washed with ethyl acetate. The combined organic layers are washed with saturated aqueous sodium bicarbonate, dried over MgSO₄, and evaporated to give a red oil. Flash chromatography (ethyl ether/hexane 2:8) afford 5.0 g (34%) of 1-(4'-t-butylidimethylsilyloxy-3'-methoxyphenyl)-1-hexene-3,5-dione, mp = 81-85°C.

Example 22B

1-(4'-Hydroxy-3'-methoxyphenyl)-6,6,6-trifluoro-1-hexene-3,5-dione

According to the procedure of Example 22A, 1-(4'-hydroxy-3'-methoxyphenyl)-1-butene-3-one is reacted with lithium di-isopropylamide (3.0 equiv) and trifluoroacetic anhydride (2.0 equiv) to afford 1-(4'-hydroxy-3'-methoxyphenyl)-6,6,6-trifluoro-1-hexene-3,5-dione in 10% yield, mp = 97-100°C.

Example 22C

1-(4'-t-Butylidimethylsilyloxy-3'-methoxyphenyl)-1-butene-3-one

To a methylene chloride (100 mL) solution of 1-(4'-hydroxy-3'-methoxyphenyl)-1-butene-3-one (10.0 g, 52 mmol) is added triethyl amine (5.8 g, 57.2 mmol) and t-butylchlorodimethyl silane (15.7 g, 104 mmol). The reaction is stirred at room temperature for 12 hours under an argon atmosphere. The mixture is washed with water (3 × 100 mL) and brine (100 mL). The organic layer is dried over MgSO₄ and evaporated. Flash chromatography (ethyl ether/hexane 1:1) affords 12.8 g (80%) of the desired 1-(4'-t-butylidimethylsilyloxy-3'-methoxyphenyl)-1-butene-3-one, mp = 65-68°C.

Example 22D

1-(4'-t-Butylidimethylsilyloxy-3',5'-dimethoxyphenyl)-1-hexene-3,5-dione

According to the procedure of Example 22C, 1-(4'-hydroxy-3',5'-dimethoxyphenyl)-1-hexene-3,5-dione is reacted with t-butylchlorodimethyl silane to afford 1-(4'-t-butylidimethylsilyloxy-3',5'-dimethoxyphenyl)-1-hexene-3,5-dione suitably pure for further use.

Example 23

5-Chloromethyl-3-trifluoromethylisoxazole

A solution of 3-trifluoromethyl-5-hydroxymethylisoxazole [(700 mg, 4.2 mmol) See K. Tanaka *et al*, *Bull. Chem. Soc. Jpn.*, **57**, 2184 (1984)] and triphenylphosphine (1.43 g, 5.4 mmol) in carbontetrachloride (50 ml) and dichloromethane (20 ml) is heated at reflux for 6 hours. The reaction mixture is then cooled to room temperature and passed through a silica gel plug. Concentration affords in 82% yield the desired 5-chloromethyl-3-trifluoromethylisoxazole of sufficient purity for the following step (Example 24).

Example 24

Isoxazole, 3-trifluoromethyl-5-methyltriphenylphosphonium chloride

Triphenylphosphine (1.016 g, 3.9 mmol) and 5-chloromethyl-3-trifluoromethylisoxazole (0.72 g, 3.9 mmol) are dissolved in toluene (40 ml) and heated at reflux for 24 hours. Upon cooling, a white precipitate formed and was filtered to afford in 60% yield the desired isoxazole, 3-trifluoromethyl-5-methyltriphenylphosphonium chloride, mp = 245-250°C.

Example 25

5-Isloxazoleethanol, α -[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]-3-methyl-

3,5-Dimethylisoxazole (26 g, 2.7×10^{-1} M) is dissolved in dry THF (100 ml) and cooled to -78°C with an acetone dry ice bath. *n*-Butyllithium (2.7×10^{-1} M) is added dropwise via a syringe. The 3,5-dimethylisoxazole, *n*-butyllithium mixture is stirred at -78°C for two hours under an argon atmosphere. 3,5-Di-*tert*-butyl-4-hydroxybenzaldehyde (18 g, 17.68×10^{-2} M) is dissolved in dry THF (200 ml) and added dropwise via cannula under an argon atmosphere to the 3,5-dimethylisoxazole and *n*-butyllithium mixture. The reaction mixture is allowed to warm to room temperature and is stirred for an additional eight hours. The reaction mixture is evaporated to near dryness, redissolved in ethyl acetate (500 ml), washed with distilled water, then brine. The organic phase is dried over sodium sulfate, filtered, evaporated to dryness, and chromatographed on flash silica using hexane:ethyl acetate 3:2 to give 16.8 g (70% yield) of the desired 5-isoxazoleethanol, α -[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]-3-methyl-, mp = $145-147^{\circ}\text{C}$. Analysis for $\text{C}_{22}\text{H}_{26}\text{O}_3\text{N}$ (331.46). Calculated C-72.47, H-8.81, N-4.22. Found C-72.31, H-8.84, N-4.24.

Example 25A5-Isloxazoleethanol, α -[4-hydroxyphenyl]-3-methyl-

According to the procedure of Example 25, 3,5-dimethylisoxazole is reacted with 4-hydroxybenzaldehyde to afford 5-isoxazoleethanol, α -[4-hydroxyphenyl]-3-methyl-, pure enough for subsequent use.

Example 25B5-Isloxazoleethanol, α -[3,5-dimethyl-4-hydroxyphenyl]-3-methyl-

According to the procedure of Example 25, 3,5-dimethylisoxazole is reacted with 3,5-dimethyl-4-hydroxybenzaldehyde to afford 5-isoxazoleethanol, α -[3,5-dimethyl-4-hydroxyphenyl]-3-methyl-, in 31% yield.

Example 25C5-Isloxazoleethanol, α -[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]-3-phenyl-

According to the procedure of Example 25, 3-phenyl-5-methylisoxazole is reacted with 3,5-bis(1,1-dimethylethyl)-4-hydroxybenzaldehyde to afford 5-isoxazoleethanol, α -[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]-3-phenyl-, in 33% yield.

Example 25D5-Isloxazoleethanol, α -[3,5-dimethoxy-4-hydroxyphenyl]-3-phenyl-

According to the procedure of Example 25, 3-phenyl-5-methylisoxazole is reacted with 3,5-dimethoxy-4-hydroxybenzaldehyde to afford 5-isoxazoleethanol, α -[3,5-dimethoxy-4-hydroxyphenyl]-3-phenyl-, in 30% yield.

Example 25E5-Isloxazoleethanol, α -[3-methoxy-4-hydroxyphenyl]-3-phenyl-

According to the procedure of Example 25, 3-phenyl-5-methylisoxazole is reacted with vanillin to afford 5-isoxazoleethanol, α -[3-methoxy-4-hydroxyphenyl]-3-phenyl-, in 41% yield.

Example 25F5-Isoxazoleethanol, α -[3,5-dibromo-4-hydroxyphenyl]-3-methyl-

5 According to the procedure of Example 25, 3,5-dimethylisoxazole is reacted with 3,5-dibromo-4-hydroxybenzaldehyde to afford 5-isoxazoleethanol, α -[3,5-dibromo-4-hydroxyphenyl]-3-methyl-, in 53% yield. Mp = 139-143°C.

Example 25G5-Isoxazoleethanol, α -[3,5-dichloro-4-hydroxyphenyl]-3-methyl-

10 According to the procedure of Example 25, 3,5-dimethylisoxazole is reacted with 3,5-dichloro-4-hydroxybenzaldehyde to afford 5-isoxazoleethanol, α -[3,5-dichloro-4-hydroxyphenyl]-3-methyl-, in 61% yield. Mp = 129-131°C.

Example 25H

20

5-Isoxazoleethanol, α -[2-hydroxy-3-methoxyphenyl]-3-methyl-

According to the procedure of Example 25, 3,5-dimethylisoxazole is reacted with 2-hydroxy-3-methoxybenzaldehyde to afford 5-isoxazoleethanol, α -[2-hydroxy-3-methoxyphenyl]-3-methyl-, as an oil in 63% yield.

Example 25J5-Isoxazoleethanol, α -[2-hydroxy-3,5-dibromophenyl]-3-methyl-

30 According to the procedure of Example 25, 3,5-dimethylisoxazole is reacted with 3,5-dibromosalicylaldehyde to afford 5-isoxazoleethanol, α -[2-hydroxy-3,5-dibromophenyl]-3-methyl-, in 47% yield.

35

Example 25K5-Isoxazoleethanol, α -[2-hydroxy-3,5-dichlorophenyl]-3-methyl-

40 According to the procedure of Example 25, 3,5-dimethylisoxazole is reacted with 3,5-dichlorosalicylaldehyde to afford 5-isoxazoleethanol, α -[2-hydroxy-3,5-dichlorophenyl]-3-methyl-, in 74% yield, mp = 151-153°C.

Example 265-Isoxazoleethanol, α -[4-hydroxy-3-methoxyphenyl]-3-methyl-

45 According to the procedure of Example 25, 3,5-dimethylisoxazole (10 g, 0.103 mole) is reacted with *n*-butyllithium (0.103 mole) and vanillin (7.8 g, 0.0515 mole) to afford 5-isoxazoleethanol, α -(4-hydroxy-3-methoxyphenyl)-3-methyl-, (10 g, 78%); mp = 128-135°C.

55

Example 27

5-Isoxazoleethanol, α -[4-hydroxy-3,5-bis(1-methylethyl)phenyl]-3-methyl-

According to the procedure of Example 25, 3,5-dimethylisoxazole (11.5 g, 0.049 mole) is reacted with *n*-butyllithium (0.049 mole) and 3,5-bis(1-methylethyl)-4-hydroxybenzaldehyde (5 g, 0.0243 mole) to afford 5-isoxazoleethanol, α -(4-hydroxy-3,5-bis(1-methylethyl)phenyl)-3-methyl-(4.0 g, 56%); mp = 79°C.

Example 285-Isoxazoleethanol, α -(4-hydroxy-3,5-dimethoxyphenyl)-3-methyl-

According to the procedure of Example 25, 3,5-dimethylisoxazole (4.26 g, 0.044 mole) is reacted with *n*-butyllithium (0.044 mole) and 4-hydroxy-3,5-dimethoxybenzaldehyde (4.00 g, 0.022 mole) to afford 5-isoxazoleethanol, α -(4-hydroxy-3,5-dimethoxyphenyl)-3-methyl-(3.82 g, 63%); mp = 142°C.

Example 295-[β -(4'-Hydroxy-3',5'-bis(1,1-dimethylethyl)phenyl) ethenyl]-3-methylisoxazole

A solution of 5-isoxazoleethanol, α -[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]-3-methyl-(16.8 g, 0.0507 mole) in HCl-saturated methanol (300 mL) is refluxed for 24 hr. The reaction mixture is evaporated to dryness, redissolved in ethyl acetate (400 mL), neutralized with 15% aqueous sodium bicarbonate and washed with brine. The residual organics are then dried (sodium sulfate) and concentrated to give a residue. Flash chromatography (hexane/ethyl acetate 3:1) affords 11.0 g of 5-[β -(4'-hydroxy-3',5'-bis(1,1-dimethylethyl)phenyl)ethenyl]-3-methylisoxazole; mp = 174-175°C. Calculated C-76.64, H, 8.68, N, 4.47. Found C-76.60; H, 8.62; N, 4.33.

Example 305-[β -(4'-Hydroxy-3'-methoxyphenyl)ethenyl]-3-methylisoxazole

According to the procedure of Example 29, 5-isoxazoleethanol, α -(4-hydroxy-3-methoxyphenyl)-3-methyl-(10 g, 0.04 mole) is reacted with HCl-methanol to afford 5-[β -(4'-hydroxy-3'-methoxyphenyl)ethenyl]-3-methylisoxazole (7.5 g, 80%); mp = 123-127°C. Calculated C-67.52, H-5.67, N-6.06. Found C-67.14, H-5.68, N-5.96.

Example 315-[β -(4'-Hydroxy-3',5'-bis(1-methylethyl)phenyl) ethenyl]-3-methylisoxazole

According to the procedure of Example 29, 5-isoxazoleethanol, α -(4-hydroxy-3,5-bis(1-methylethyl)phenyl)-3-methyl-(4.0 g, 0.013 mole) is reacted with HCl-methanol to afford 5-[β -(4'-hydroxy-3',5'-bis(1-methylethyl)phenyl)ethenyl]-3-methylisoxazole (3.9 g, 98%); mp = 119-120°C. Calculated C-75.76, H-8.12, N-4.91. Found C-75.68, H-8.16, N-4.96.

Example 325-[β -(4'-Hydroxy-3',5'-dimethoxyphenyl)ethenyl]-3-methylisoxazole

According to the procedure of Example 29, 5-isoxazoleethanol, α -(4-hydroxy-3,5-dimethoxyphenyl)-3-methyl-(3.00 g, 10.7 mmole) is reacted with HCl-methanol to afford 5-[β -(4'-hydroxy-3',5'-dimethoxyphenyl)ethenyl]-3-methylisoxazole (1.92 g, 69%); mp = 152-157°C. Calculated C-64.36, H-5.79, N-5.36. Found C-64.30, H-5.74, N-5.34.

Example 32A5-[β -(4'-Hydroxyphenyl)ethenyl]-3-methylisoxazole

- 6 According to the procedure of Example 29, 5-isoxazoleethanol, α -[4-hydroxyphenyl]-3-methyl- is reacted with HCl-methanol to afford 5-[β -(4'-hydroxyphenyl)ethenyl]-3-methylisoxazole (6.74 g, 14%); mp = 117-119°C.

10 Example 32B5-[β -(4'-Hydroxy-3',5'-dimethylphenyl)ethenyl]-3-methylisoxazole

- 15 According to the procedure of Example 29, 5-isoxazoleethanol, α -[3,5-dimethyl-4-hydroxyphenyl]-3-methyl- is reacted with HCl-methanol to afford 5-[β -(4'-hydroxy-3',5'-dimethylphenyl)ethenyl]-3-methylisoxazole (0.62 g, 85%); mp = 158-161°C. Analysis for $C_{14}H_{15}NO_2$. Calculated C-73.34, H-6.59, N-6.10. Found C-73.35, H-6.44, N-6.05.

20 Example 32C5-[β -(4'-Hydroxy-3',5'-bis(1,1-dimethylethyl)phenyl)ethenyl]-3-phenylisoxazole

- 25 According to the procedure of Example 29, 5-isoxazoleethanol, α -[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]-3-phenyl- is reacted with HCl-methanol to afford 5-[β -(4'-hydroxy-3',5'-bis(1,1-dimethylethyl)-phenyl) ethenyl]-3-phenylisoxazole (0.35 g, 90%); mp = 157-161°C. analysis for $C_{28}H_{29}NO_2$. Calculated C-79.96, H-7.78, N-3.73. Found C-79.78, H-7.74, N-3.71.

30 Example 32D5-[β -(4'-Hydroxy-3',5'-dimethoxyphenyl)ethenyl]-3-phenylisoxazole

- 35 According to the procedure of Example 29, 5-isoxazoleethanol, α -[3,5-dimethoxy-4-hydroxyphenyl]-3-phenyl- is reacted with HCl-methanol to afford 5-[β -(4'-hydroxy-3',5'-dimethoxyphenyl) ethenyl]-3-phenylisoxazole (1.1 g, 95%); mp = 135-140°C. Analysis for $C_{19}H_{17}NO_4$. Calculated C-70.58, H-5.30, N-4.33. Found C-70.70, H-5.17, N-4.25.

40 Example 32E5-[β -(4'-Hydroxy-3'-methoxyphenyl)ethenyl]-3-phenylisoxazole

- 45 According to the procedure of Example 29, 5-isoxazoleethanol, α -[3-methoxy-4-hydroxyphenyl]-3-phenyl- is reacted with HCl-methanol to afford 5-[β -(4'-hydroxy-3'-methoxyphenyl)ethenyl]-3-phenylisoxazole (1.0 g, 95%) mp = 148-152°C. Analysis for $C_{18}H_{15}NO_3$. Calculated C-73.71, H-5.15, N-4.77. Found C-73.33, H-5.13, N-4.69.

50 Example 32F5-[β -(2'-Hydroxy-3'-methoxyphenyl)ethenyl]-3-methylisoxazole

- 55 According to the procedure of Example 29, 5-isoxazoleethanol, α -[2-hydroxy-3-methoxyphenyl]-3-methyl- is reacted with HCl-methanol to afford 5-[β -(2'-hydroxy-3'-methoxyphenyl)ethenyl]-3-methylisoxazole (31%); mp = 129-131°C. Analysis for $C_{13}H_{13}NO_3$. Calculated C-67.51, H-5.68, N-6.06. Found C-67.51, H-5.70, N-5.95.

Example 32G5-[β -(4'-Hydroxy-3',5'-dibromophenyl)ethenyl]-3-methylisoxazole

- 5 According to the procedure of Example 29, 5-isoxazoleethanol, α -[3,5-dibromo-4-hydroxyphenyl]-3-methyl- is reacted with HCl-methanol to afford 5-[β -(4'-hydroxy-3',5'-dibromophenyl)ethenyl]-3-methylisoxazole (15%), mp = 167-169°C. Analysis for $C_{12}H_9Br_2NO_2$. Calculated C-40.12, H-2.53, N-3.90, Br-44.49. Found C-40.18, H-2.48, N-3.64, Br-44.88.

10

Example 32H5-[β -(4'-Hydroxy-3',5'-dichlorophenyl)ethenyl]-3-methylisoxazole

- 15 A toluene (100 mL) solution of 5-isoxazoleethanol, α -[3,5-dichloro-4-hydroxyphenyl]-3-methyl-(5.5 g, 19.1 mmol) containing a catalytic amount of p-toluenesulfonic acid is warmed to reflux for 3 hours, with azeotropic removal of water. The solvent is evaporated, and the residue is chromatographed (ethyl acetate/chloroform, 1:9) to afford 5-[β -(4'-hydroxy-3',5'-dichlorophenyl)ethenyl]-3-methylisoxazole (2.3 g, 45%); mp = 184-186°C. Analysis for $C_{12}H_9Cl_2NO_2$. Calculated C-53.35, H-3.37, N-5.19. Found C-52.98, H-3.22, N-5.28.

20

Example 32I5-[β -(2'-Hydroxy-3',5'-dibromophenyl)ethenyl]-3-methylisoxazole

- 25 According to the procedure of Example 32H, 5-isoxazoleethanol, α -[2-hydroxy-3,5-dibromophenyl]-3-methyl- is reacted with p-toluenesulfonic acid/toluene to afford 5-[β -(2'-hydroxy-3',5'-dibromophenyl)ethenyl]-3-methylisoxazole in 70% yield, mp = 150-157°C. Analysis for $C_{12}H_9NO_2Br_2$. Calculated C-40.15, H-2.53, N-3.90. Found C-39.85, H-2.47, N-3.75.

30

Example 32J5-[β -(2'-Hydroxy-3',5'-dichlorophenyl)ethenyl]-3-methylisoxazole

- 35 According to the procedure of Example 32H, 5-isoxazoleethanol, α -[2-hydroxy-3,5-dichlorophenyl]-3-methyl- is reacted with p-toluenesulfonic acid/toluene to afford 5-[β -(2'-hydroxy-3',5'-dichlorophenyl)ethenyl]-3-methylisoxazole in 80% yield, mp = 143-145°C. Analysis for $C_{12}H_9NO_2Cl_2$. Calculated C-53.36, H-3.36, N-5.19, Cl-26.25. Found C-53.12, H-3.05, N-5.11, Cl-26.26.

40

Example 334-Acetoxy-3-methoxycinnamaldehyde

- 45 Acetoxy vanillin (5.0 g, 2.6×10^{-2} M) is dissolved in dry THF (300 ml) with formyl methylenetriphenylphosphorane (78 g, 2.6×10^{-2} M) and refluxed under an atmosphere of argon for 80 hours. The reaction mixture is concentrated to dryness and chromatographed on flash silica to afford 4-acetoxy-3-methoxycinnamaldehyde (3 g, 54% yield); mp = 83-86°C.

50

Example 34

55

4-Acetoxy-3-methoxycinnamaldehyde oxime

4-Acetoxy-3-methoxycinnamaldehyde (2.0 g, 9.1×10^{-3} M) is dissolved in methanol (100 ml) and cooled to 0°C with an ice/water bath. Hydroxylamine hydrochloride (0.8 g, 1.2×10^{-2} M) is added with sodium acetate (0.97 g, 1.2×10^{-2} M) and stirred for one hour. The reaction mixture is concentrated to near dryness, and redissolved in ethyl acetate (150 ml). The residual organics are washed twice with water dried over sodium sulfate, filtered, evaporated to dryness, and chromatographed on flash silica using hexane:ethyl acetate 3:1 to afford 1.45 g of 4-acetoxy-3-methoxycinnamaldehyde oxime in 68% yield; mp = 72-74°C.

Example 353-[β -(4'-Acetoxy-3'-methoxyphenyl)ethenyl]-5-phenylisoxazole

4-Acetoxy-3-methoxycinnamaldehyde oxime (0.6, 2.5×10^{-3} M) is dissolved in dry DMF (15 ml) at 0°C under an atmosphere of argon. N-chlorosuccinimide (0.4 g, 3.0×10^{-3} M) is dissolved in dry DMF (10 ml) and transferred to the 4-acetoxycinnamaldehyde/DMF solution via cannula under an atmosphere of argon. The reaction is stirred at 0°C for nearly two hours. Phenylacetylene (1.3 g, 1.3×10^{-2} M) is mixed with triethylamine (0.38 g, 3.8×10^{-3} M) and dry DMF (5 ml). This mixture is then added via syringe to the cinnamaldehyde/NCS mixture. The entire mixture is allowed to reach room temperature and is stirred overnight. The reaction mixture is redissolved in ethyl acetate (150 ml), washed first with water, then with brine. The organics are dried over sodium sulfate, filtered, and evaporated to dryness. The residue is chromatographed on flash silica using hexane:ethyl acetate 3:2 to afford 0.45 g of 3-[β -(4'-acetoxy-3'-methoxyphenyl)ethenyl]-5-phenylisoxazole in 54% yield; mp = 139-144°C.

Example 35A3-[β -(4'-Acetoxy-3'-methoxyphenyl)ethenyl]-5-(2'-hydroxyethyl)isoxazole

According to the procedure of Example 35, 4-acetoxy-3-methoxycinnamaldehyde oxime is reacted with N-chlorosuccinimide and 3-buten-1-ol to afford 3-[β -(4'-acetoxy-3'-methoxyphenyl)ethenyl]-5-(2'-hydroxyethyl)isoxazole (2.0 g, 50%); mp = 62-64°C.

Example 35B3-[β -(4'-Acetoxy-3'-methoxyphenyl)ethenyl]-5-hydroxymethylisoxazole

According to the procedure of Example 35, 4-acetoxy-3-methoxycinnamaldehyde oxime is reacted with N-chlorosuccinimide and propargyl alcohol to afford 3-[β -(4'-acetoxy-3'-methoxyphenyl)ethenyl]-5-hydroxymethylisoxazole (2.5 g, 67%); sufficiently pure for further use.

Example 35C3-Phenyl-5-methylisoxazole

Benzaldehyde oxime (20.0 g, 0.165 mole) is dissolved in DMF (250 mL) and cooled to 4°C in an ice bath. N-chlorosuccinimide (33.1 g, 0.248 mole) is added to the reaction mixture, and the mixture is stirred for one hour. Isopropenyl acetate (41.3 g, 0.41 mole) and triethyl amine (24.8 g, 0.247 mole) are added via an equal-pressure addition funnel to the solution of benzaldehyde hydroxamic chloride. The ice bath is removed and the mixture is stirred for 24 hours. The mixture is then taken up into ethyl acetate (1 L) and washed with water (2 \times 1 L). The organic layer is dried over sodium sulfate and evaporated to give a residue. Flash chromatography (hexane/ethyl acetate 9:1) affords 12.0 g of 3-phenyl-5-methylisoxazole as a yellow oil.

Example 363-[β -(4'-Hydroxy-3'-methoxyphenyl)ethenyl]-5-phenylisoxazole

- 5 3-[β -(4'-Acetoxy-3'-methoxyphenyl)ethenyl]-5-phenylisoxazole (0.3 g, 8.9×10^{-4} M) is dissolved in MeOH (20 ml) and stirred with sodium methoxide (0.1 g, 1.8×10^{-3} M) for one hour. The reaction mixture is concentrated to dryness, redissolved in ethyl acetate (75 ml), and washed twice with water. The organics are dried over sodium sulfate, filtered, evaporated to dryness, and chromatographed on flash silica using hexane:ethyl acetate to afford 0.25 g of 3-[β -(4'-hydroxy-3'-methoxyphenyl)ethenyl]-5-phenylisoxazole in
 10 97% yield; mp = 135-138°C. Calculated C-73.71, H-5.15, N-4.77. Found C-73.35, H-5.35, N-4.77.

Example 36A3-[β -(4'-Hydroxy-3'-methoxyphenyl)ethenyl]-5-(2'-hydroxyethyl)isoxazole

- According to the procedure of Example 36, 3-[β -(4'-acetoxy-3'-methoxyphenyl)ethenyl]-5-(2'-hydroxyethyl)isoxazole is reacted with sodium methoxide/methanol to afford 3-[β -(4'-hydroxy-3'-methoxyphenyl)ethenyl]-5-(2'-hydroxyethyl)isoxazole (0.24 g, 93%); mp = 127-131°C. Analysis for $C_{14}H_{15}NO_4$. Calculated C-
 20 64.35, H-5.99, N-5.36. Found C-64.40, H-5.79, N-5.15.

Example 36B3-[β -(4'-Hydroxy-3'-methoxyphenyl)ethenyl]-5-hydroxymethylisoxazole

- According to the procedure of Example 36, 3-[β -(4'-acetoxy-3'-methoxyphenyl)ethenyl]-5-hydroxymethylisoxazole is reacted with sodium methoxide/methanol to afford 3-[β -(4'-hydroxy-3'-methoxyphenyl)ethenyl]-5-hydroxymethylisoxazole (0.6 g, 79%); mp = 139-143°C. Analysis for $C_{13}H_{13}NO_4 \cdot 0.4 H_2O$. Cal-
 30 culated C-63.36, H-5.46, N-5.50. Found C-63.34, H-5.53, N-5.30.

Example 373-[β -(4'-Acetoxy-3'-methoxyphenyl)ethenyl]-5-methylpyrazole

- To a solution of 3-[β -(4'-hydroxy-3'-methoxyphenyl)ethenyl]-5-methylpyrazole (2.50 g, 10.9 mmols) in pyridine (30 mL) is added acetic anhydride (5.54 g, 54.3 mmols). The reaction is stirred at room temperature for 12 hr. After this time, the reaction mixture is added dropwise to vigorously stirred ice-water
 40 (500 mL). The resulting suspension is allowed to stand for 90 minutes and the crude diacetyl product (3.2 g) is collected by filtration. The crude diacetyl product is dissolved in acetone (300 mL), water (30 mL) and methanol (30 mL). Basic alumina (100 g, Woelm basic, tic grade) is added and the reaction mixture refluxed for 4.5 hr. The alumina is filtered and rinsed with acetone (100 mL). The combined filtrate is evaporated and the residue partitioned between water and ethyl acetate. The organic layer is dried ($MgSO_4$) and con-
 45 centrated to afford, after flash chromatography [chloroform:ethyl acetate 3:1], 1.71 g of pure 3-[β -(4'-acetoxy-3'-methoxyphenyl)ethenyl]-5-methylpyrazole; mp = 131-136°C. Analysis for $C_{15}H_{15}N_2O_3$. Calculated C-66.16, H-5.92, N-10.29. Found C-66.49, H-5.92, N-10.29.

Example 383-[β -(4'-Hydroxy-3',5'-bis(1,1-dimethylethyl)phenyl)ethenyl]-5-methylpyrazole

- A solution of 5-[β -(4'-hydroxy-3',5'-bis(1,1-dimethylethyl)phenyl)ethenyl]-3-methylisoxazole (2.00 g, 6.4 mmols), water (0.115 g, 6.4 mmols) and molybdenum hexacarbonyl (1.27 g, 48 mmols) in acetonitrile (75 mL) is refluxed for 12 hr under a nitrogen atmosphere. The reaction mixture is cooled and evaporated to dryness. The residue is dissolved in methanol (250 mL) and acidified to pH = 1 with 4 N HCl. After stirring for 4 hr at room temperature, the methanol is evaporated. The resulting aqueous solution is neutralized with

1 N NaOH and the organic material is extracted into ethyl acetate. The organic solution is passed through a pad of silica gel (150 g) and further eluted with chloroform (400 mL). The combined filtrate is evaporated to dryness, taken up in a minimal amount of ethyl acetate, and passed through a silica gel pad again. The resulting crude diketone (1.25 g) is then suspended in a mixture of acetic acid (100 mL) and 97% hydrazine (1.0 mL). The reaction mixture is stirred for 12 hr at room temperature. Evaporation affords a gummy residue which solidifies upon stirring in water (100 mL) for 30 minutes. The solid is purified by flash chromatography [methylenedichloride/ethyl acetate 1:1] to afford 1.10 g of 3- β -(4'-hydroxy-3',5'-bis(1,1-dimethylethyl)phenyl)ethenyl]-5-methylpyrazole; mp = 218-223°C. Analysis for $C_{20}H_{28}N_2O$. Calculated C-76.88, H-9.03, N-8.96. Found C-76.63, H-8.84, N-9.02.

Example 38A

3- β -(4'-Hydroxy-3',5'-bis(1-methylethyl)phenyl)ethenyl]-5-methylpyrazole

According to the procedure of Example 38, 5- β -(4'-hydroxy-3',5'-bis(1-methylethyl)phenyl)ethenyl]-3-methylisoxazole is reacted with molybdenum hexacarbonyl to afford 3- β -(4'-hydroxy-3',5'-bis(1,1-dimethylethyl)phenyl)ethenyl]-5-methylpyrazole in 53% yield, mp = 183-184°C. Analysis for $C_{18}H_{24}N_2O \cdot 0.1 H_2O$. Calculated C-75.54, H-8.52, N-9.79. Found C-75.33, H-8.72, N-9.46.

Example 38B

3- β -(4'-Hydroxy-3',5'-dimethoxyphenyl)ethenyl]-5-methylpyrazole

According to the procedure of Example 38, 5- β -(4'-hydroxy-3',5'-dimethoxyphenyl)ethenyl]-3-methylisoxazole is reacted with molybdenum hexacarbonyl to afford 3- β -(4'-hydroxy-3',5'-dimethoxyphenyl)ethenyl]-5-methylpyrazole in 56% yield, mp = 153-158°C. Analysis for $C_{14}H_{16}N_2O_3$. Calculated C-64.60, H-6.19, N-10.76. Found C-64.36, H-6.24, N-10.61.

Example 38C

3- β -(4'-Hydroxy-3'-methoxyphenyl)ethenyl]-5-methylpyrazole

According to the procedure of Example 38, 5- β -(4'-hydroxy-3'-methoxyphenyl)ethenyl]-3-methylisoxazole is reacted with molybdenum hexacarbonyl to afford 3- β -(4'-hydroxy-3'-methoxyphenyl)ethenyl]-5-methylpyrazole in 85% yield, mp = 145-147°C. Analysis for $C_{13}H_{14}N_2O_2$. Calculated C-67.81, H-6.13, N-12.17. Found C-67.62, H-6.18, N-11.93.

Example 38D

3- β -(2'-Hydroxy-3',5'-dibromophenyl)ethenyl]-5-methylpyrazole

According to the procedure of Example 38, 5- β -(2'-hydroxy-3',5'-dibromophenyl)ethenyl]-3-methylisoxazole is reacted with molybdenum hexacarbonyl to afford 3- β -(2'-hydroxy-3',5'-dibromophenyl)ethenyl]-5-methylpyrazole in 30% yield. mp = 191-200°C. Analysis for $C_{12}H_{10}N_2OBr_2$. Calculated C-40.26, H-2.82, N-7.82, Br-44.64. Found C-40.02, H-2.82, N-7.71, Br-44.35.

Example 39

5-[β -(4'-Hydroxy-3',5'-dimethoxyphenyl)ethenyl]-3-methylisoxazole

A solution of 3-methyl-5-isoxazoleacetic acid (10.0 g, 71 mmol) [see reference R. G. Micetich, Canadian J. Chem., **48**, 2006 (1970)], syringaldehyde (12.9 g, 71 mmol), piperidine (0.6 g, 7 mmol), and acetic acid (0.42 g, 7 mmol) in toluene (500 mL) is refluxed with azeotropic removal of water for 3 hours. The mixture is cooled, and the solid is collected by filtration. The solid is dissolved in pyridine (100 mL) and refluxed for 4 hours. The solvent is evaporated, and the residue purified by flash chromatography (methanol/chloroform 1:9) and recrystallization (ethyl acetate) to afford 5-[β -(4'-hydroxy-3',5'-dimethoxyphenyl)ethenyl]-3-methylisoxazole (7.73 g, 42%); mp = 152-155°C. Analysis for $C_{14}H_{15}NO_4$. Calculated C-64.36, H-5.79, N-5.36. Found C-64.16, H-5.69, N-5.28.

Example 39A5-[β -(3'-Methoxy-4'-hydroxy-5'-bromophenyl)ethenyl]-3-methylisoxazole

According to the procedure of Example 39, 5-bromovanillin is reacted with 3-methyl-5-isoxazoleacetic acid to afford 5-[β -(3'-methoxy-4'-hydroxy-5'-bromophenyl)ethenyl]-3-methylisoxazole in 55% yield, mp = 178-180°C. Analysis for $C_{13}H_{12}NO_3Br$. Calculated C-50.33, H-3.90, N-4.54. Found C-50.35, H-3.88, N-4.44.

Example 405-[α -Carbomethoxy- β -(4'-hydroxy-3',5'-dimethoxy phenyl)ethenyl]-3-methylisoxazole

A solution of 3-methyl-5-isoxazoleacetic acid methyl ester (7.75 g, 50 mmol), syringaldehyde (9.1 g, 50 mmol), piperidine (0.42 g, 5 mmol), and acetic acid (0.3 g, 5 mmol) in toluene (250 mL) is refluxed for 12 hours with azeotropic removal of water. Concentration affords a residue which is taken up in ethyl acetate and adsorbed onto dry silica gel pad. The pad is washed with ethyl acetate, and the filtrate is then evaporated. Recrystallization from ethyl acetate/hexane gives 5-[α -carbomethoxy- β -(4'-hydroxy-3',5'-dimethoxyphenyl)ethenyl]-3-methylisoxazole (7.9 g, 50%), mp = 130-132°C. Analysis for $C_{16}H_{17}NO_6$. Calculated C-60.17, H-5.38, N-4.39. Found C-60.17, H-5.53, N-4.34.

Example 415-[α -Carbomethoxy- β -(4'-hydroxy-3,5-dichlorophenyl) ethenyl]-3-methylisoxazole

According to the procedure of Example 40, 3-methyl-5-isoxazoleacetic acid methyl ester is reacted with 3,5-dichloro-4-hydroxybenzaldehyde to afford 5-[α -carbomethoxy- β -(4'-hydroxy-3',5'-dichlorophenyl)ethenyl]-3-methylisoxazole (37%), mp = 198-200°C. Analysis for $C_{14}H_{11}Cl_2NO_4$. Calculated C-51.24, H-3.39, N-4.27, Cl-21.61. Found C-50.99, H-3.28, N-4.37, Cl-20.88.

Example 425-[α -Carbomethoxy- β -(4'-hydroxy-3',5'-bis(1-methylethyl)phenyl)ethenyl]-3-methylisoxazole

According to the procedure of Example 40, 3-methyl-5-isoxazoleacetic acid methyl ester is reacted with 4-hydroxy-3,5-bis(1-methylethyl) benzaldehyde to afford 5-[α -carbomethoxy- β -(4'-hydroxy-3',5'-bis(1,1-dimethylmethyl)phenyl)ethenyl]-3-methylisoxazole (41%), mp = 126-128°C. Analysis for $C_{26}H_{25}NO_4$. Calculated C-69.94, H-7.35, N-4.08. Found C-69.88, H-7.29, N-3.72.

Example 43

5-[α -Carbomethoxy- β -(4'-hydroxy-3',5'-dimethyl)phenyl)ethenyl]-3-methylisoxazole

According to the procedure of Example 40, 3-methyl-5-isoxazoleacetic acid methyl ester is reacted with 4-hydroxy-3,5-dimethylbenzaldehyde to afford 5-[α -carbomethoxy- β -(4'-hydroxy-3',5'-dimethylphenyl)ethenyl]-3-methylisoxazole (45%), mp = 176-178°C. Analysis for $C_{16}H_{17}NO_4$. Calculated C-66.88, H-5.98, N-4.88. found C-66.46, H-6.01, N-4.85.

Example 445-[α -Carbomethoxy- β -(4'-hydroxy-3',5'-dibromophenyl) ethenyl]-3-methylisoxazole

According to the procedure of Example 40, 3-methyl-5-isoxazoleacetic acid methyl ester is reacted with 4-hydroxy-3,5-dibromobenzaldehyde to afford 5-[α -carbomethoxy- β -(4'-hydroxy-3',5'-dibromophenyl)ethenyl]-3-methylisoxazole (48%), mp = 164-166°C. Analysis for $C_{14}H_{11}Br_2NO_4$. Calculated C-40.32, H-2.66, N-3.36, Br-38.32. Found C-40.34, H-2.61, N-3.30, Br-38.38.

Example 455-[α -Carbomethoxy- β -(4'-hydroxy-3'-methoxyphenyl) ethenyl]-3-methylisoxazole

According to the procedure of Example 40, 3-methyl-5-isoxazoleacetic acid methyl ester is reacted with vanillin to afford 5-[α -carbomethoxy- β -(4'-hydroxy-3'-methoxyphenyl)ethenyl]-3-methylisoxazole (51%), mp = 117-119°C. Analysis for $C_{15}H_{15}NO_5$. Calculated C-60.17, H-5.38, N-4.39. Found C-60.17, H-5.53, N-4.34.

Example 465-[α -Carbomethoxy- β -(4'-hydroxy-3',5'-bis(1,1-dimethylethyl)phenyl)ethenyl]-3-methylisoxazole

According to the procedure of Example 40, 3-methyl-5-isoxazoleacetic acid methyl ester is reacted with 4-hydroxy-3,5-bis(1,1-dimethylethyl) benzaldehyde (with the exception of a longer reaction time of 6 days) to afford 5-[α -carbomethoxy- β -(4'-hydroxy-3',5'-bis(1,1-dimethylethyl)phenyl) ethenyl]-3-methylisoxazole (25%), mp = 131-136°C. Analysis for $C_{20}H_{23}NO_4$. Calculated C-71.13, H-7.87, N-3.77. Found C-71.41, H-7.99, N-3.75.

Example 46A5-[α -Carbomethoxy- β -(3'-methoxy-4'-hydroxy-5'-bromophenyl)ethenyl]-3-methylisoxazole

According to the procedure of Example 40, 3-methyl-5-isoxazoleacetic acid methyl ester is reacted with 5-bromovanillin to afford 5-[α -carbomethoxy- β -(3'-methoxy-4'-hydroxy-5'-bromophenyl)ethenyl]-3-methylisoxazole in 38% yield, mp = 138-140°C. Analysis for $C_{15}H_{14}NO_5Br$. Calculated C-48.93, H-3.84, N-3.80, Br-21.70. Found C-49.10, H-3.88, N-3.96, Br-21.35.

Example 475-[α -Carboxy- β -(4'-hydroxy-3',5'-dimethoxyphenyl) ethenyl]-3-methylisoxazole

A solution of 5-[α -carbomethoxy- β -(4'-hydroxy-3',5'-dimethoxyphenyl)ethenyl]-3-methylisoxazole (3.0 g, 9.4 mmol) in 1 N aqueous KOH (100 mL) is stirred at room temperature for 18 hours. The solution is acidified with aqueous HCl and the precipitate formed is collected by filtration. Recrystallization from methanol/water 1:9 affords 5-[α -carboxy- β -(4'-hydroxy-3',5'-dimethoxyphenyl)ethenyl]-3-methylisoxazole (2.17 g, 76%); mp = 217-219°C. Analysis for $C_{15}H_{15}NO_6$. Calculated C-59.01, H-4.96, N-4.59. Found C-58.96, H-4.90, N-4.51.

Example 485-[α -Carboxy- β -(4'-hydroxy-3',5'-dichlorophenyl) ethenyl]-3-methylisoxazole

5 According to the procedure of Example 47, 5-[α -Carbomethoxy- β -(4'-hydroxy-3',5'-dichlorophenyl) ethenyl]-3-methylisoxazole is saponified to afford 5-[α -carboxy- β -(4'-hydroxy-3',5'-dichlorophenyl) ethenyl]-3-methylisoxazole (64%), mp = 229-230°C. Analysis for $C_{13}H_9Cl_2NO_4$. Calculated C-49.70, H-2.89, N-4.46, Cl-22.57. Found C-49.46, H-2.65, N-4.17, Cl-22.54.

10

Example 495-[α -Carboxy- β -(4'-hydroxy-3',5'-bis(1-methylethyl)phenyl)ethenyl]-3-methylisoxazole

15 According to the procedure of Example 47, 5-[α -carbomethoxy- β -(4'-hydroxy-3',5'-bis(1-methylethyl)-phenyl)ethenyl]-3-methylisoxazole is saponified to afford 5-[α -carboxy- β -(4'-hydroxy-3',5'-bis(1,1-dimethyl-methyl)phenyl)ethenyl]-3-methylisoxazole (83%), mp = 192-194°C. Analysis for $C_{26}H_{22}NO_4$. Calculated C-69.27, H-7.05, N-4.25. Found C-69.51, H-7.06, N-4.16.

20

Example 505-[α -Carboxy- β -(4'-hydroxy-3'-methoxyphenyl)ethenyl]-3-methylisoxazole

25 According to the procedure of Example 47, 5-[α -carbomethoxy- β -(4'-hydroxy-3'-methoxyphenyl) ethenyl]-3-methylisoxazole is saponified to afford 5-[α -carboxy- β -(4'-hydroxy-3'-methoxyphenyl)ethenyl]-3-methylisoxazole (73%), mp = 198-199°C. Analysis for $C_{14}H_{13}NO_5$. Calculated C-61.08, H-4.77, N-5.09. Found C-60.74, H-4.72, N-4.65.

30

Example 515-[α -Carboxy- β -(4'-hydroxy-3',5'-bis(1,1-dimethylethyl)phenyl)ethenyl]-3-methylisoxazole

35 According to the procedure of Example 47, 5-[α -carbomethoxy- β -(4'-hydroxy-3',5'-bis(1,1-dimethylethyl)-phenyl)ethenyl]-3-methylisoxazole is saponified to afford 5-[α -carboxy- β -(4'-hydroxy-3',5'-bis(1,1-dimethylethyl)phenyl)ethenyl]-3-methylisoxazole (85%), mp = 219-221°C. Analysis for $C_{24}H_{27}NO_4$. Calculated C-70.55, H-7.63, N-3.92. Found C-70.48, H-7.64, N-3.84.

40

Example 525-[α -Carboxy- β -(4'-hydroxy-3',5'-dimethylphenyl) ethenyl]-3-methylisoxazole

45 According to the procedure of Example 47, 5-[α -carbomethoxy- β -(4'-hydroxy-3',5'-dimethylphenyl)-ethenyl]-3-methylisoxazole is saponified to afford 5-[α -carboxy- β -(4'-hydroxy-3',5'-dimethylphenyl)ethenyl]-3-methylisoxazole (90%), mp 228-229°C. Analysis for $C_{16}H_{15}NO_4$. Calculated C-65.91, H-5.54, N-5.13. Found C-66.30, H-5.62, N-5.06.

50

Example 535-[α -Carboxy- β -(4'-hydroxy-3',5'-dibromophenyl) ethenyl]-3-methylisoxazole

55 According to the procedure of Example 47, 5-[α -carboxy- β -(4'-hydroxy-3',5'-dibromophenyl) ethenyl]-3-methylisoxazole is saponified to afford 5-[α -carboxy- β -(4'-hydroxy-3',5'-dibromophenyl) ethenyl]-3-methylisoxazole (77%), mp = 229-230°C. Analysis for $C_{13}H_9Br_2NO_4$. Calculated C-38.74, H-2.02, N-3.48, Br-39.65. Found C-38.75, H-2.20, N-3.45, Br-39.93.

Example 53A5-[α -Carboxy- β -(3'-methoxy-4'-hydroxy-5'-bromophenyl)ethenyl]-3-methylisoxazole

- 5 According to the procedure of Example 47, 5-[α -carboxy- β -(3'-methoxy-4'-hydroxy-5'-bromophenyl)-ethenyl]-3-methylisoxazole is saponified to afford 5-[α -carboxy- β -(3'-methoxy-4'-hydroxy-5'-bromophenyl)-ethenyl]-3-methylisoxazole (98%), mp = 225-227°C. Analysis for $C_{14}H_{12}NO_5Br$. Calculated C-47.47, H-3.42, N-3.96, Br-22.56. Found C-47.45, H-3.42, N-4.14, Br-22.84.

10

Example 543-[β -(4'-Hydroxy-3',5'-bis(1,1-dimethylethyl)phenyl)ethenyl]-5-methylisoxazole

- 15 According to the procedure of Example 39, 5-methyl-3-isoxazoleacetic acid is reacted with 4-hydroxy-3,5-bis(1,1-dimethylethyl)benzaldehyde to afford 3-[β -(4'-hydroxy-3',5'-bis(1,1-dimethylethyl)phenyl)ethenyl]-5-methylisoxazole in 44% yield. Mp = 172-174°C. Analysis for $C_{20}H_{22}NO_2$. Calculated C-76.64, H-8.68, N-4.47. Found C-76.44, H-8.58, N-4.59.

20

Example 54A3-[β -(4'-Hydroxy-3',5'-dibromophenyl)ethenyl]-5-methylisoxazole

- 25 According to the procedure of Example 39, 5-methyl-3-isoxazoleacetic acid is reacted with 4-hydroxy-3,5-dibromobenzaldehyde to afford 3-[β -(4'-hydroxy-3',5'-dibromophenyl)ethenyl]-5-methylisoxazole in 85% yield, mp = 178-182°C. Analysis for $C_{12}H_9NO_2Br_2$. Calculated C-40.14, H-2.53, N-3.90. Found C-40.09, H-2.54, N-3.99.

30

Example 553-[β -(4'-Hydroxy-3',5'-bis(1,1-dimethylethyl)phenyl)ethenyl]-5-carboxymethylisoxazole

- 35 A solution of 3-[β -(4'-hydroxy-3',5'-bis(1,1-dimethylethyl)phenyl)ethenyl]-5-methylisoxazole (0.5 g, 1.6 mmol) in THF (15 mL) is cooled to -78°C. N-butyllithium (2 mL, 3.2 mmol) is added dropwise, and the solution is stirred for an additional 20 minutes. The reaction mixture is then poured onto dry ice and allowed to come to room temperature under a flow of argon gas. Water (100 mL) is added and the mixture is extracted with ethyl acetate (200 mL). The organic layer is washed with saturated aqueous NH_4Cl (1X) and
40 water (2X), then dried over sodium sulfate. Chromatography of the residue on silica (chloroform/methanol 95:5 containing 0.1% acetic acid) affords 3-[β -(4'-hydroxy-3',5'-bis(1,1-dimethylethyl)phenyl)ethenyl]-5-carboxymethylisoxazole (0.4 g, 70%), mp = 205-210°C. Analysis for $C_{21}H_{27}NO_4$. Calculated C-70.56, H-7.61, N-3.92. Found C-70.19, H-7.86, N-3.80.

45

Example 533-[β -(4'-Hydroxy-3',5'-bis(1,1-dimethylethyl)phenyl)ethenyl]-5-methylpyrazole

- 50 A solution of 3-methyl-5-pyrazoleacetic acid (20.0 g, 0.14 mol), 4-hydroxy-3,5-bis(1,1-dimethylethyl)benzaldehyde (33.4 g, 0.14 mol), piperidine (1.2 g, 0.014 mol), and acetic acid (0.85 g, 0.014 mol) in toluene (1 L) is refluxed for 24 hours with azeotropic removal of water. The mixture is cooled and evaporated to afford a residue, which is dissolved in methanol/ethyl acetate (1:9) and extracted with water (300 mL). The organic layer is evaporated and recrystallized from hexane/ethyl acetate (5:1) to afford 3-[β -(4'-hydroxy-3',5'-bis(1,1-dimethylethyl)phenyl)ethenyl]-5-methylpyrazole (27.4 g, 61%). Mp = 220-224°C.
55 Analysis for $C_{20}H_{22}N_2O$. Calculated C-76.88, H-9.03, N-8.96. Found C-77.15, H-9.14, N-9.09.

Example 573-[β -(4'-Hydroxy-3',5'-dimethylphenyl)ethenyl]-5-methylpyrazole

- 5 According to the procedure of Example 56, 3-methyl-5-pyrazoleacetic acid is reacted with 4-hydroxy-3,5-dimethylbenzaldehyde to afford 3-[β -(4'-hydroxy-3',5'-dimethylphenyl)ethenyl]-5-methylpyrazole in 50% yield. Mp = 185-188°C. Analysis for $C_{14}H_{16}N_2O$. Calculated C-73.66, H-7.06, N-12.27. Found C-73.68, H-7.13, N-12.26.

10

Example 583-[β -(4'-Hydroxy-3',5'-dibromophenyl)ethenyl]-5-methylpyrazole

- 15 According to the procedure of Example 56, 3-methyl-5-pyrazoleacetic acid is reacted with 4-hydroxy-3,5-dibromobenzaldehyde to afford 3-[β -(4'-hydroxy-3',5'-dibromophenyl)ethenyl]-5-methylpyrazole in 51% yield. Mp = 101-108°C. Analysis for $C_{12}H_{10}Br_2N_2O$. Calculated C-40.26, H-2.82, N-7.82. Found C-39.92, H-2.83, N-7.70.

20

Example 58A3-[β -(3'-Methoxy-4'-hydroxy-5'-bromophenyl)ethenyl]-5-methylpyrazole

- 25 According to the procedure of Example 56, 3-methyl-5-pyrazoleacetic acid is reacted with 3-methoxy-4-hydroxy-5-bromobenzaldehyde to afford 3-[β -(3'-methoxy-4'-hydroxy-5'-bromophenyl)ethenyl]-5-methylpyrazole in 20% yield; mp = 158-160°C. Analysis for $C_{13}H_{13}N_2O_2Br$. Calculated C-50.50, H-4.24, N-9.06. Found C-50.84, H-4.37, N-8.88.

30

Example 593-[α -Carbomethoxy- β -(4'-hydroxy-3',5'-dibromophenyl)ethenyl]-5-methylpyrazole

- 35 According to the procedure of Example 40, 3-methyl-5-pyrazoleacetic acid methyl ester is reacted with 4-hydroxy-3,5-dibromobenzaldehyde to afford 3-[α -carbomethoxy- β -(4'-hydroxy-3',5'-dibromophenyl)ethenyl]-5-methylpyrazole in 40% yield, mp = 182-184°C.

40 Example 59A3-[α -Carbomethoxy- β -(4'-hydroxy-3',5'-dichlorophenyl)ethenyl]-5-methylpyrazole

- 45 According to the procedure of Example 40, 3-methyl-5-isoxazoleacetic acid methyl ester is reacted with 4-hydroxy-3,5-dichlorobenzaldehyde to afford 3-[α -carbomethoxy- β -(4'-hydroxy-3',5'-dichlorophenyl)ethenyl]-5-methylpyrazole in 16% yield, mp = 210-212°C. Analysis for $C_{14}H_{12}N_2O_3Cl_2$. Calculated C-51.39, H-3.70, N-8.56. Found C-51.77, H-3.80, N-8.43.

50 Example 59B3-[α -Carbomethoxy- β -(3'-methoxy-4'-hydroxy-5'-bromophenyl)ethenyl]-5-methylpyrazole

- 55 According to the procedure of Example 40, 3-methyl-5-pyrazoleacetic acid methyl ester is reacted with 5-bromovanillin to afford 3-[α -carbomethoxy- β -(3'-methoxy-4'-hydroxy-5'-bromophenyl)ethenyl]-5-methylpyrazole in 17% yield, mp = 200-201°C. Analysis for $C_{15}H_{15}N_2O_4Br$. Calculated C-49.06, H-4.13, N-7.63. Found C-49.29, H-4.11, N-7.66.

Example 603-Methyl-5-pyrazoleacetic acid methyl ester

5 A solution of 3-methyl-5-pyrazoleacetic acid (5.0 g, 35.7 mmol) [see C. Ainsworth, J. Amer. Chem. Soc., 76, 3172 (1954)] in methanol (50 mL) is saturated with HCl gas. The mixture is warmed to reflux for 5 hours, after which the solution is concentrated. The solid residue is taken up in chloroform and washed with 0.5 N aqueous sodium bicarbonate, dried over MgSO_4 , and evaporated to afford 3-methyl-5-pyrazoleacetic acid methyl ester (3.9 g, 70%) sufficiently pure for further use.

Example 613-[α -Carboxy- β -(4'-hydroxy-3',5'-dibromophenyl)ethenyl]-5-methylpyrazole

15 According to the procedure of Example 47, 3-[α -carboxy- β -(4'-hydroxy-3',5'-dibromophenyl)ethenyl]-5-methylpyrazole is saponified to afford 3-[α -carboxy- β -(4'-hydroxy-3',5'-dibromophenyl)ethenyl]-5-methylpyrazole in 88% yield, mp = 194-197°C. Analysis for $\text{C}_{13}\text{H}_{10}\text{N}_2\text{O}_3\text{Br}_2$. Calculated C-38.83, H-2.51, N-6.97. Found C-38.80, H-2.51, N-6.98.

Example 623-[α -Carboxy- β -(4'-hydroxy-3',5'-dichlorophenyl)ethenyl]-5-methylpyrazole

25 According to the procedure of Example 47, 3-[α -carboxy- β -(4'-hydroxy-3',5'-dichlorophenyl)ethenyl]-5-methylpyrazole is saponified to afford 3-[α -carboxy- β -(4'-hydroxy-3',5'-dichlorophenyl)ethenyl]-5-methylpyrazole in 71% yield, mp = 223-225°C. Analysis for $\text{C}_{13}\text{H}_{10}\text{N}_2\text{O}_3\text{Cl}_2$. Calculated C-49.87, H-3.23, N-8.95, Cl-22.64. Found C-49.66, H-3.13, N-8.82, Cl-22.72.

Example 633-[α -Carboxy- β -(3'-methoxy-4'-hydroxy-5'-bromophenyl)ethenyl]-5-methylpyrazole

35 According to the procedure of Example 47, 3-[α -carboxy- β -(3'-methoxy-4'-hydroxy-5'-bromophenyl)ethenyl]-5-methylpyrazole is saponified to afford 3-[α -carboxy- β -(3'-methoxy-4'-hydroxy-5'-bromophenyl)ethenyl]-5-methylpyrazole in 65% yield, mp = 145-148°C. Analysis for $\text{C}_{14}\text{H}_{13}\text{N}_2\text{O}_4\text{Br} \cdot 0.5 \text{ H}_2\text{O}$. Calculated C-46.42, H-3.62, N-7.74, Br-22.06. Found C-46.64, H-3.82, N-7.82, Br-22.01.

Example 645-[β -(4'-Hydroxy-3'-methoxyphenyl)ethenyl]-3-methylisothiazole

45 To a suspension of 4-acetoxy-3-methoxybenzyltriphenyl phosphonium chloride [L. Lonsky et al., Manatshefte für chemie, 107, 685-695 (1976)] (2.24 g, 4.7 mmol) in dry THF (100 mL) and DMSO (4 mL) at 0°C under an argon atmosphere is added sodium hydride, 60% suspension in oil (0.18 g, 4.7 mmol). The reaction mixture is stirred at room temperature for one hour and then 3-methylisothiazole-5-carboxaldehyde (D. Buttmore et al., JCS, 2063, (1963)) (0.6 g, 4.7 mmol) is added. The reaction mixture is stirred at room temperature for 18 hours. The reaction mixture is poured into a saturated solution of ammonium chloride (250 mL) and the product is extracted into ethyl acetate (2x200 mL). The crude intermediate 5-[β -(4'-acetoxy-3'-methoxyphenyl)ethenyl]-3-methylisothiazole (mixture of cis and trans isomers) is purified by flash chromatography (silica, 25% EtOAc/ CH_2Cl_2). The intermediate acetoxy compound (0.72 g) is dissolved in methanol (20 mL) and treated with sodium methoxide (0.26 g) and the reaction mixture is stirred at room temperature for one hour. The reaction mixture is diluted with ethyl acetate (50 mL) and washed with water (50 mL). The organic layer is dried (MgSO_4) and evaporated to give 5-[β -(4'-hydroxy-3'-methoxyphenyl)-

ethenyl]-3-methylisothiazole (0.54 g). Flash chromatography (silica, 5% EtOAc/CHCl₃) gave separation of the trans and cis isomers.

High R_f isomer (0.35 g, 30%), R_f = 0.53 (silica, 8:1/CH₂Cl₂:EtOAc); mp = 187-189°C.

Low R_f isomer (0.10 g, 9%), R_f = 0.43 (silica, 8:1/CH₂Cl₂:EtOAc); mp = 120-124°C.

- 5 The usefulness of the compounds of the present invention as inhibitors of lipoxigenase enzyme or other related biochemical actions is demonstrated by their effectiveness in various standard test procedures. A description of each procedure follows.

10 5-Lipoxygenase Assay Using Isolated Human Leukocytes (5LOA₂)

The formation of 5-HETE in human leukocytes is considered a measure of 5-lipoxygenase activity. The protocol is described in the following.

- 15 Fresh heparinized or EDTA treated human blood is mixed with 6% dextran-3% dextrose in isotonic saline in the ratio 0.25 ml dextran solution per 1.0 ml blood. After mixing the blood is allowed to sit at room temperature for about 90 minutes while the RBC's settle. During this period, the plasma is removed with a plastic pipette to nalgens tubes.

- 20 The plasma is centrifuged at 800 rpm (125 kg) on the Beckman Td-b refrigerated centrifuge to remove the platelets (which remain in the supernatant). The pellet, consisting of leukocytes and erythrocytes, is treated with 10 ml 0.87% ammonium chloride at room temperature for four minutes, lysing the red cells. At the end of four minutes the cells are diluted with a 2^x volume of phosphate buffered saline, pH 7.4, and centrifuged for ten minutes. The cells are washed three times with the phosphate buffered saline. Any of the pelleted cell matter which is not easily resuspended is discarded during the washings--the material contains platelets (12-lipoxygenase activity).

- 25 After washing, the cells are resuspended in phosphate buffered saline containing 1.0 mM calcium and 0.5 mM magnesium. After counting, the cells are diluted to 1.5-2.0 × 10⁷ leukocytes per milliliter.

To each polypropylene reaction tube is added 0.48 ml leukocytes in Ca-Mg phosphate buffered saline, pH 7.4; 1-5 μl test compound dissolved in DMSO and buffer; or DMSO for control tubes.

The tubes preincubate at 37°C for five minutes.

- 30 The reaction is started by adding 20 μl of the following, 0.5 μl, 20 mM arachidonic acid--final concentration = 20 μM; 1 μl, 5 mM calcium ionophore A23187--final concentration = 10 μM; and 18.5 μl buffer.

- 35 The reaction proceeds for five minutes, then is stopped by adding 0.5 ml, 0.5 mM ice-cold Tris buffer, pH 8.0. The tubes are chilled on ice for ten minutes and then extracted three times with a total of 3.5 ml ethyl acetate (3.0 ml removed).

The tubes can be stored at this point. For extended storage, the tubes should be filled with nitrogen.

The ethyl acetate is evaporated with a Sorvall Speed-Vac. The residue is dissolved in ethanol. The tubes can also be stored at this point at -20°C under nitrogen.

A portion of the ethanol solution is injected into the HPLC system for 5-HETE quantitation.

- 40 The HPLC system consists of Hewlett-Packard 1040A UV spectrophotometry system with an HP85 computer. Injections are made automatically with a Waters WISP 710B. The pump is a Spectra Physics SP8700. Peaks are measured with a Hewlett-Packard 3390A integrator. An RP C-18 column is used. The solvent system is isocratic; the solvent is 70% methanol and 30% 0.01M sodium acetate, pH 5.7, pumped at 1.0 ml/min. The flow is monitored at 235 nm for 5-HETE quantitation. Using a 15 cm Alltech Nucleosil C-18 5 μM column provides for a sample turnaround time of about 16 minutes.

IC₅₀ is calculated as the amount of test agent that causes 50% inhibition of the formation of 5-HETE relative to the control.

50 Cyclooxygenase Enzyme Assay

- Additionally, inhibition of cyclooxygenase is considered a measure of relevance to the pathophysiology of the above noted diseases. For example, see "Inhibition of Immunoglobulin E-Mediated, Antigen-Induced Monkey Asthma and Skin Reactions by 5,8,11,14-Eicosatetraynoic Acid," by Roy Patterson, M.D. and Kathleen E. Harris, B.S. in, J. Allergy Clin. Immunol., Vol. 67, No. 2, pp. 146-152.

The assay consists of incubating 2 mg bovine seminal vesicle powder with 2 mM epinephrine, 2.5 mM reduced glutathione, 100 μ M arachidonic acid, and the test agent for 20 minutes. The reaction mixture is acidified and extracted with ethyl acetate (3 \times 1.0 ml) and the pooled extract is evaporated to dryness using a Speed Vac Concentrator or under a stream of nitrogen. The residue was dissolved in ethanol. An aliquot is applied on 20 \times 20 cm silica gel plate and developed using water:ethyl acetate:hexane:acetic acid (60:54:25:12.5, upper phase) to separate PGE₂ from arachidonic acid. ¹⁴C-PGE₂ formed is identified by co-chromatography with authentic ³H-PGE₂ and the amount of radioactivity is quantitated using an automatic TLC linear scanner (Berthold, Pittsburgh, Pennsylvania) linked to an Apple II-e computer and an IC₅₀ is calculated as the amount of test compound causing 50% inhibition of cyclooxygenase enzyme relative to the control.

The above defined value for each of tested compounds of the present invention having the noted Example numbers are as found in the following Table 2.

TABLE 2

<u>Example #</u>	<u>5-LO(IC₅₀ in μM)</u>	<u>BSV(IC₅₀ in μM)</u>
1	1.0	66.0
2	53.0	67.0
3	15.0	44.0
4	4.3	40.0
5	7.1	33.0
6	>20	-
7	5.5	34.0
8	>20	-
9	7.5	56.0
10	18.0	-
12	29.0	31.0
29	7.8	10.0
30	6.8	-
32	1.8	-

ARBL/ARBC Whole Cell 5-Lipoxygenase and Cyclooxygenase Assays

Materials

The rat basophilic leukemia cell line (RBL-1) was obtained from the American Type Culture Collection (Rockville, MD).

Radioimmuno assay (RIA) kits of LTB₄ and PGF_{2 α} were obtained from Amersham (Arlington Heights, IL) and Seragen (Boston, MA) respectively.

All tissue culture media were obtained from GIBCO (Grand Island, NY).

Method

RBL-1 cells were grown in culture (suspension) in Eagle's minimum essential medium supplemented with 12% fetal bovine serum and 1:100 antibiotic-antimycotic mixture at 37°C in an incubator supplied with air-5% carbon dioxide. Cells were harvested by centrifugation. They were washed with cold phosphate buffered saline pH 7.4 (PBS; NaCl, 7.1 g; Na₂HPO₄, 1.15 g; KH₂PO₄, 0.2 g; and KCl, 0.2 g/L). Cells were finally suspended in PBS containing 0.88 μM calcium at a density of 2.4 × 10⁶ cells/ml. Cells were incubated with and without test agent (in DMSO) (1% DMSO was without effect on arachidonic acid metabolism) for ten minutes at room temperature. Calcium ionophore A23187 (5 μM) was added and cells were incubated for seven minutes at 37°C. The reaction was stopped by chilling the tubes in ice for ten minutes. Cells were separated by centrifugation and the supernatant was stored at -20°. Aliquots (100 μl) were analyzed for LTB₄ and PGF_{2α} using radioimmuno assay kits as provided by the supplier.

Table 3 contains biochemical data obtained from this RBL-1 whole cell assay.

15

20

25

30

35

40

45

50

55

TABLE 3

5

	<u>Example #</u>	<u>ARBL (IC₅₀, μM)^a</u>	<u>ARBC (IC₅₀, μM)^b</u>
10	1	1.6	2.5
	5	0.8	13.0
	7	9.5	6.8
	10	5.2	11.0
15	10A	3.8	4.5
	10D	0.3	>30
	10G-1	4.5	>30
20	10G-2	4.1	>30
	10H	9.5	>30
	10I	>10	>25
	10J	>10	30.0
25	10K	>20	-
	10L	>20	-
	10M	>30	>30
30	10N	>20	>30
	10P	12.0	>30
	10Q	>30	>30
35	29	2.4	1.5
	30	5.5	12.0
	31	1.0	10.0
	32	1.4	22.0
40	32B	4.5	1.2
	32D	<1.0	5.4
	32E	4.5	2.8
45	32F	12.0	30.0
	32G	11.0	16.0
	32H	21.0	13.8
	32J	4.7	>30
50	36	-	-
	36A	21.5	N
	37	3.5	-
55	38	3.0	0.9

TABLE 3 (Continued)

	<u>Example #</u>	<u>ARBL (IC₅₀, μM)^a</u>	<u>ARBC (IC₅₀, μM)^b</u>
5	38A	3.6	14.0
	38B	2.6	>30
10	39A	4.0	9.3
	40	>10	>25
	41	>30	5.0
15	43	17.0	2.2
	44	12.0	5.2
	46	10.0	2.0
	46A	>30	12.0
20	51	20.0	2.5
	52	>20	>20
	54	2.7	\approx 0.3
25	54B	3.3	4.2
	55	17.0	>30
	57	0.82	4.6
30	58	5.7	14.5

^aRBL-1 intact cell 5-lipoxygenase IC₅₀ values (μ M)

^bRBL-1 intact cell cyclooxygenase IC₅₀ values (μ M)

35 Accordingly, the present invention also includes a pharmaceutical composition for treating one of the above diseases or conditions comprising an antidisease or anticondition effective amount of a compound of the formula I as defined above together with a pharmaceutically acceptable carrier.

The present invention further includes a method of use of a compound of the formula I for the manufacturing of pharmaceutical compositions for the treatment of the above named diseases.

40 For preparing pharmaceutical compositions from the compounds described by this invention, inert, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, dispersible granules, capsules, cachets, and suppositories. A solid carrier can be one or more substances which may also act as diluents, flavoring agents, solubilizers, lubricants, suspending agents, binders or tablet disintegrating agents; it can also be encapsulating material. In powders, the carrier is a
45 finely divided solid which is in admixture with the finely divided active compound. In the tablet the active compound is mixed with carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain from 5 or 10 to about 70 percent of the active ingredient. Suitable solid carriers are magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. The term "preparation" is intended to include the
50 formulation of the active compound with encapsulating material as carrier providing a capsule in which the active component (with or without other carriers) is surrounded by carrier, which is thus in association with it. Similarly, cachets are included. Tablets, powders, cachets, and capsules can be used as solid dosage forms suitable for oral administration.

55 For preparing suppositories, a low melting wax such as a mixture of fatty acid glycerides or cocoa butter is first melted, and the active ingredient is dispersed homogeneously therein as by stirring. The molten homogeneous mixture is then poured into convenient sized molds, allowed to cool and thereby to solidify.

Liquid form preparations include solutions, suspensions, and emulsions. As an example may be mentioned water or water-propylene glycol solutions for parenteral injection. Liquid preparations can also be formulated in solution in aqueous polyethylene glycol solution. Aqueous solutions suitable for oral use can be prepared by dissolving the active component in water and adding suitable colorants, flavors, stabilizing and thickening agents as desired. Aqueous suspensions suitable for oral use can be made by dispersing the finely divided active component in water with viscous material, i.e., natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, and other well-known suspending agents.

Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations for either oral or parenteral administration. Such liquid forms include solutions, suspensions, and emulsions. These particular solid form preparations are most conveniently provided in unit dose form and as such are used to provide a single liquid dosage unit. Alternately, sufficient solid may be provided so that after conversion to liquid form, multiple individual liquid doses may be obtained by measuring predetermined volumes of the liquid form preparation as with a syringe, teaspoon, or other volumetric container. When multiple liquid doses are so prepared, it is preferred to maintain the unused portion of said liquid doses at low temperature (i.e., under refrigeration) in order to retard possible decomposition. The solid form preparations intended to be converted to liquid form may contain, in addition to the active material, flavorants, colorants, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like. The liquid utilized for preparing the liquid form preparation may be water, isotonic water, ethanol, glycerine, propylene glycol, and the like as well as mixtures thereof. Naturally, the liquid utilized will be chosen with regard to the route of administration, for example, liquid preparations containing large amounts of ethanol are not suitable for parenteral use.

Preferably, the pharmaceutical preparation is in unit dosage form. In such form, the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, for example, packeted tablets, capsules, and powders in vials or ampoules. The unit dosage form can also be a capsule, cachet, or tablet itself or it can be the appropriate number of any of these in packaged form.

The quantity of active compound in a unit dose of preparation may be varied or adjusted from 1 mg to 500 mg preferably to 1 to 50 mg according to the particular application and the potency of the active ingredient. The composition can, if desired, also contain other compatible therapeutic agents.

In therapeutic use as described above, the dosages may be varied depending upon the requirements of the patient, the severity of the condition being treated, and the compound being employed. Determination of the proper dosage for a particular situation is within the skill of the art. Generally, treatment is initiated with the smaller dosages which are less than the optimum dose of the compound. Thereafter the dosage is increased by small increments until the optimum effect under the circumstances is reached. For convenience, the total daily dosage may be divided and administered in portions during the day if desired.

As used herein cardiovascular diseases or conditions particularly include 1) reductions of the extent of infarct damage in a myocardial infarction, 2) prevention of recurrent myocardial infarction, 3) stroke, 4) anaphylactic shock, and 5) vasospastic disease.

An additional advantageous benefit of the cytoprotective property of the compounds of formula I are for use, for example to protect against damage from various gastro-intestinal tract conditions.

Various assays that are generally accepted can be used to measure cytoprotective ability.

The magnitude of a prophylactic or therapeutic dose of a compound of formula I will, of course, vary with the nature of the severity of the condition to be treated and with the particular compound of formula I and its route of administration. In general, the daily dose range for anti-asthmatic, anti-allergic or anti-inflammatory use and, generally uses other than cytoprotection, lies within the range of from about 10 μ g to about 20 mg per kg body weight of a mammal, preferably from about 50 μ g to about 20 mg per kg of body weight of a mammal, and most preferably from about 100 μ g to about 10 mg per kg of body weight of a mammal.

The exact amount of a compound of the formula I to be used as a cytoprotective agent will depend on, inter alia, whether it is being administered to heal damaged cells or to avoid future damage, on the nature of the damaged cells (e.g., gastro-intestinal ulcerations vs. nephrotic necrosis), and on the nature of the causative agent. An example of the use of a compound of the formula I in avoiding future damage would be co-administration of a compound of the formula I with a non-steroidal anti-inflammatory drug (for example, indomethacin) that might otherwise cause such damage. For such use, the compound of formula I is administered from 30 minutes prior up to 30 minutes after administration of the NSAID. Preferably, it is administered prior to or simultaneously with the NSAID (e.g. as a combination dosage form).

The effective daily dosage level for compounds of formula I inducing cytoprotection in mammals, especially humans, will generally range from about 0.002 mg/kg to about 100 mg/kg, preferably from about 0.02 mg/kg to about 30 mg/kg. The dosage may be administered in single or divided individual doses.

Thus, in addition to the compounds of formula I, the pharmaceutical compositions can also contain other active ingredients, such as cyclooxygenase inhibitors non-steroidal anti-inflammatory drugs (NSAIDs), peripheral analgesic agents such as zomepirac, diflunisal and the like. The weight ratio of the compound of the formula I to the second active ingredient may be varied and will depend upon the effective dose of each ingredient. Generally, an effective dose of each will be used. Thus, for example, when a compound of the formula I is combined with an NSAID, the weight ratio of the compound of the formula I to the NSAID will generally range from about 1000:1 to about 1:1000, preferably about 200:1 to about 1:200. Combinations of a compound of the formula I and other active ingredients will generally also be within the aforementioned range, but in each case, an effective dose of each active ingredient should be used.

Combinations of a compound of the formula I and other active ingredients will generally be in the aforementioned ratios.

NSAIDs can be characterized into five groups:

- (1) the propionic acid derivatives;
- (2) the acetic acid derivatives;
- (3) the fenamic acid derivatives;
- (4) the biphenylcarboxylic acid derivatives; and
- (5) the oxicams or a pharmaceutically acceptable salt thereof.

The propionic acid derivatives which may be used comprise: ibuprofen, ibuprofen aluminum, indoprofen, ketoprofen, naproxen, benoxaprofen, flurbiprofen, fenoprofen, fenbufen, pirofen, carprofen, oxaprozin, pranoprofen, miroprofen, tiaprofen, alminoprofen, tiaprofen, fluprofen and buclocic acid. Structurally related propionic acid derivatives having similar analgesic and anti-inflammatory properties are also intended to be included in this group.


Thus, "propionic acid derivatives" as defined herein are non-narcotic analgesics/non-steroidal anti-inflammatory drugs having a free $-\text{CH}(\text{CH}_3)\text{COOH}$ or $-\text{CH}_2\text{CH}_2\text{COOH}$ group (which optionally can be in the form of a pharmaceutically acceptable salt group, e.g., $-\text{CH}(\text{CH}_3)\text{COO}^-\text{Na}^+$ or $-\text{CH}_2\text{CH}_2\text{COO}^-\text{Na}^+$), typically attached directly or via a carbonyl function to a ring system, preferably to an aromatic ring system.

The acetic acid derivatives which may be used comprise: indomethacin, which is a preferred NSAID, sulindac, tolmetin, zomepirac, diclofenac, fenclofenac, alclofenac, ibufenac, isoxepac, furofenac, tiopinac, zidometacin, acemetacin, fentiazac, clidanac, oxpinac, and fenclozic acid. Structurally related acetic acid derivatives having similar analgesic and anti-inflammatory properties are also intended to be encompassed by this group.

Thus, "acetic acid derivatives" as defined herein are non-narcotic analgesics/non-steroidal anti-inflammatory drugs having a free $-\text{CH}_2\text{COOH}$ group (which optionally can be in the form of a pharmaceutically acceptable salt group, e.g., $-\text{CH}_2\text{COO}^-\text{Na}^+$), typically attached directly to a ring system, preferably to an aromatic or heteroaromatic ring system.

The fenamic acid derivatives which may be used comprise: mefenamic acid, meclofenamic acid, flufenamic acid, niflumic acid and tolfenamic acid. Structurally related fenamic acid derivatives having similar analgesic and anti-inflammatory properties are also intended to be encompassed by this group.


Thus, "fenamic acid derivative" as defined herein are non-narcotic analgesics/non-steroidal anti-inflammatory drugs which contain the basic structure:



which can bear a variety of substituents and in which the free $-\text{COOH}$ group can be in the form of a pharmaceutically acceptable salt group, e.g., $-\text{COO}^-\text{Na}^+$.

The biphenylcarboxylic acid derivatives which can be used comprise: diflunisal and flufenisal. Structurally related biphenylcarboxylic acid derivatives having similar analgesic and anti-inflammatory properties are also intended to be encompassed by this group.


Thus, "biphenylcarboxylic acid derivatives" as defined herein are non-narcotic analgesics/non-steroidal anti-inflammatory drugs which contain the basic structure:



which can bear a variety of substituents and in which the free -COOH group can be in the form of a pharmaceutically acceptable salt group, e.g., -COO-Na⁺.

The oxicams which can be used in the present invention comprise: piroxicam, sudoxicam, isoxicam and 4-hydroxyl-1,2-benzothiazine 1,1-dioxide 4-(N-phenyl)-carboxamide. Structurally related oxicams having similar analgesic and anti-inflammatory properties are also intended to be encompassed by this group.

Thus, "oxicams" as defined herein are non-narcotic analgesics/non-steroidal anti-inflammatory drugs which have the general formula:



wherein R is an aryl or heteroaryl ring system.


The following NSAIDs may also be used: acemetacin, alminoprofen, amfenac sodium, aminoprofen, anitrazafen, antrafenine, auranofin, bendazac lysinate, benzydamine, beprozin, properamole, bufezolac, carprofen, cinmetacin, ciproquazone, clidanac, cloximate, dazidamine, deboxamet, delmetacin, detomidine, dexindoprofen, diacerein, di-fisalamine, difenpyramide, emorfazone, enfenamic acid, enolicam, epirizole, etersalate, etodolac, etofenamate, fanetizole mesylate, fenclofenac, fenclorac, fendosal, fenflumizole, fentiazac, feprazone, floctafenine, flunixin, flunoxaprofen, fluproquazone, fopirtoline, fosfosal, furcloprofen, furofenac, glucametacin, guimesal, ibuprofen, isofezolac, isonixim, isoprofen, isoxepac, isoxicam, lefetamine HCl, leflunomide, lofemizole, lonazolac calcium, lotifazole, loxoprofen, lysin, clonixinate, meclofenamate sodium, meseclazone, microprofen, nabumetone, nictindole, nimesulide, orpanoxin, oxametacin, oxapadol, oxaprozin, perisoxal citrate, pimeprofen, pimmetacin, pirofen, pirazolac, pirfenidone, pirprofen, pranoprofen, proglumetacin maleate, proquazone, pyridoxiprofen, sudoxicam, suprofen, talmetacin, talniflumate, tenoxicam, thiazolinobutazone, thielavin B, tiaprofenic acid, tiaramide HCl, tiflamizole, timegadine, tiroxaprofen, tolfenamic acid, tolpadol, tryptamid, ufenamate, and zidometacin.

Finally, NSAIDs which may also be used include the salicylates, specifically aspirin, and the phenylbutazones, and pharmaceutically acceptable salts thereof.

Pharmaceutical compositions comprising the formula I compounds may also contain as the second active ingredient, antihistaminic agents such as benadryl, dramamine, histadyl, phenergan and the like. Alternatively, they may include prostaglandin antagonists such as those disclosed in European Patent Application 11,067 or thromboxane antagonists such as those disclosed in U.S. 4,237,160. They may also contain histidine decarboxylase inhibitors such as α -fluoromethylhistidine, described in U.S. 4,325,961. The compounds of the formula I may also be advantageously combined with an H₁ or H₂-receptor antagonist, such as for instance cimetidine, ranitidine, terfenadine, famotidine, temelastine, acrivastine, loratadine, cetirizine, taziflyline, azelastine, aminothiadiazoles disclosed in EP 81102976.8 and like compounds, such as those disclosed in U.S. Patent Nos. 4,283,408; 4,362,736; 4,394,508 and European Patent Application No. 40,696. The pharmaceutical compositions may also contain a K⁺/H⁺ ATPase inhibitor such as omeprazole, disclosed in U.S. Pat. 4,255,431, and the like. Each of the references referred to in this paragraph is hereby incorporated herein by reference.

Claims

1. A compound of the formula (I)




and pharmaceutically acceptable salts thereof; wherein

(1) -- is a single or double bond;


15 (2) R, R₁, and R₂ may be the same or different and are selected from the group consisting of hydrogen, lower alkyl, hydroxy, OR₃ wherein R₃ is lower alkyl, C(O)OR₄ wherein R₄ is hydrogen or lower alkyl, OC(O)R₃ wherein R₃ is independently as defined above, C(O)R₃ wherein R₃ is independently as defined above, NR₆R₇, wherein R₆ and R₇ may be the same or different and are hydrogen or lower alkyl, NHC(O)R₃ wherein R₃ is independently as defined above, NHCHO, NHSO₂R₃ wherein R₃ is independently as defined above, NHCONHR₄ wherein R₄ is as defined above, hydroxymethyl, halogen, trifluoromethyl, SR₄ wherein R₄ is independently as defined above, or nitro;

20 (3) Q is (CH₂)_n wherein n is an integer of zero to four, CH=CH or




wherein R₄ is independently as defined above;

30 (4) X and Y are (i) N, (ii) NR₅ wherein R₅ is hydrogen, lower alkyl,




35 wherein R₁['] and R₂['] may be the same or different and are hydrogen or lower alkyl, C(O)R₄ wherein R₄ is independently as defined above, cycloalkyl of from three to twenty carbons having of from three to eight ring carbons, aryl, or aralkyl, (iii) O, (iv) S; with the proviso that X and Y cannot both be N, NR₅, O or S at once and with the proviso that one of X and Y cannot be O at the same time the other of X and Y is S or NR₅ and that one of X and Y cannot be S at the same time the other of X and Y is NR₅;

40 (5) Z is H, lower alkyl, aryl, aralkyl, OC(O)R₃ wherein R₃ is independently as defined above, C(O)OR₄ wherein R₄ is independently as defined above, C(O)R₃ wherein R₃ is independently as defined above, CH-(R₁['])CO₂R₂['] wherein R₁['] and R₂['] are independently as defined above, halogen, trifluoromethyl,



wherein R, R₁ and R₂ are independently as defined above, heteroaryl, or heteroaralkyl; with the overall proviso that when one of R, R₁, and R₂ is 2-hydroxy, X is O, Y is N and Q is CH=CH, then Z cannot be H or alkyl, and also with the overall proviso that when R, R₁, and R₂ are hydroxy or lower alkyl, Y and X are N or NH, n is zero then Z cannot be furyl or phenyl unsubstituted or substituted with halogen, trifluoromethyl, alkyl, alkoxy or NO₂.

2. A compound of Claim 1 wherein Q is (CH₂)_n wherein n is zero or two, CH=CH, or



wherein R_4 is as defined above, and wherein X and Y are N and NR_5 with R_5 as defined above, NR_5 and N, O and N or N and O, S and N or N and S.

3. 3- $[\beta$ -(4'-hydroxy-3'-methoxyphenyl)ethenyl]-5-methylpyrazole or

3- $[\beta$ -(4'-acetoxy-3'-methoxyphenyl)ethenyl]-5-methylpyrazole

4. 5- $[\beta$ -(4'-hydroxy-3',-5'-bis(1,1-dimethylethyl)phenyl)ethenyl]-3-methylisoxazole or

5- $[\beta$ -(4'-hydroxy-3',5'-dimethoxyphenyl)ethenyl]-3-methylisoxazole.

5. 3- $[\beta$ -(4'-hydroxy-3',5'-bis(1,1-dimethylethyl)phenyl)ethenyl]-5-methylpyrazole.

6. A pharmaceutical composition for use as inhibitors of 5-lipoxygenase or cyclooxygenase comprising a 5-lipoxygenase or cyclooxygenase inhibiting amount of a compound of formula (I) according to claim 1.


7. A sunscreen composition for use as a sunscreen comprising an ultraviolet absorbing amount of a compound of Claim 1 not including the overall provisos and a suitable carrier therefor.

8. A pharmaceutical composition as claimed in Claim 1 additionally comprising an effective amount of a second active ingredient that is a non-steroidal anti-inflammatory drug; a peripheral analgesic agent; a cyclooxygenase inhibitor; a leukotriene antagonist; an antihistaminic agent; a prostaglandin antagonist or a thromboxane antagonist.


9. A method of use of a compound of general formula I according to claim 1 alone or in combination with other pharmaceutically active compounds for the manufacture of pharmaceutical compositions for treating inflammation, arthritis, ulcers, allergies, asthma, psoriasis and cardiovascular conditions in mammals suffering therefrom or for the manufacture of compositions for protection from ultraviolet light.

10. A process for the preparation of a compound of formula I of Claim 1 which comprises

(1) reacting a compound of the formula




wherein R, R_1 , and R_2 are as defined above; and a compound of the formula




wherein $\text{--}\text{--}$ and Z is as defined above, Y and X are O and N, N and O, S and N or N and S, and Ph is phenyl, in the presence of a suitable base; to obtain the compound of formula I wherein Z is as defined above, Y and X are O and N, N and O, S and N, N and S, and R, R_1 , and R_2 are as defined above;

(2) treating a compound of the formula




wherein R, R_1 , R_2 , and Z are as defined above and Q is $\text{HC}=\text{CH}$ or $(\text{CH}_2)_n$; with hydrazine, and optionally hydrogenating the $\text{--}\text{--}$ bond when it is a double bond; to obtain a compound of formula I wherein X and Y are N and NH, and R, R_1 , R_2 , and Z are as defined above;

(3) (step a) treating a compound of the formula



wherein R, R₁, and R₂ are as defined above; with a compound of the formula




wherein Li is lithium, W is H or CO₂R₄ wherein R₄ is as defined above and Z is as defined above; and then (step b) treating the product of the step a with hydrochloric acid in methanol; to obtain a product of formula I wherein X and Y are O and N and Q is CH=C,

$$\downarrow$$


and R, R₁, R₂, and Z are as defined above;

and, alternatively, further treating the compound obtained in step b with Mo(CO)₆ and then hydrazine to obtain the compound of the formula I wherein X and Y are N and NH and R, R₁, R₂, or Z are as defined above;

(4) treating a compound of the formula



wherein R, R₁, and R₂ are as defined above, with N-chlorosuccinimide and base followed by treating with a compound of the formula




to obtain the compound of formula I wherein X is N and Y is O, Q is HC≡CH and R, R₁, R₂, and Z are as defined above.


(5a) treating a compound of the formula I of Claim 1 wherein one of X and Y is oxygen and the other is nitrogen with acetonitrile in the presence of molybdenum hexacarbonyl; and then

b) treating with hydrazine to obtain a compound of the formula I of Claim 1 wherein X and Y are nitrogen

(6a) contacting a compound of the formula




wherein R, R₁, and R₂ are as defined above and a compound of the formula




wherein Z and R₄ are as defined above, and one of X and Y is NH and other is N, or one of X and Y is O and the other is N, and then in situ either

b) dehydrating and decarboxylating the product of step (a) wherein R₄ is hydrogen to obtain a compound of the formula



wherein one of X and Y is NH and the other is N or one of X and Y is O and the other is N, and R, R₁, R₂ and Z are as defined above; or


c) dehydrating the product of step (a) (in situ) wherein R₄ is alkyl to obtain the compound of formula



wherein one of X and Y is NH and the other is N, or one of X and Y is O and the other is N, and R, R₁, R₂ and R₄ are as defined above.

Claims for the following contracting states: AT, ES, GR

1. A process for the preparation of a compound of the formula (I)




I

and pharmaceutically acceptable salts thereof; wherein

(1) --- is a single or double bond;


(2) R, R₁, and R₂ may be the same or different and are selected from the group consisting of hydrogen, lower alkyl, hydroxy, OR₃ wherein R₃ is lower alkyl, C(O)OR₄ wherein R₄ is hydrogen or lower alkyl, OC(O)R₃ wherein R₃ is independently as defined above, C(O)R₃ wherein R₃ is independently as defined above, NR₆R₇, wherein R₆ and R₇ may be the same or different and are hydrogen or lower alkyl, NHC(O)R₃ wherein R₃ is independently as defined above, NHCHO, NHSO₂R₃ wherein R₃ is independently as defined above, NHCONHR₄ wherein R₄ is as defined above, hydroxymethyl, halogen, trifluoromethyl, SR₄ wherein R₄ is independently as defined above, or nitro;

(3) Q is (CH₂)_n wherein n is an integer of zero to four, CH=CH or




wherein R₄ is independently as defined above;

(4) X and Y are (i) N, (ii) NR₅ wherein R₅ is hydrogen, lower alkyl,




wherein R₁' and R₂' may be the same or different and are hydrogen or lower alkyl, C(O)R₄ wherein R₄ is independently as defined above, cycloalkyl of from three to twenty carbons having of from three to eight ring carbons, aryl, or aralkyl, (iii) O, (iv) S; with the proviso that X and Y cannot both be N, NR₅, O or S at once and with the proviso that one of X and Y cannot be O at the same time the other of X and Y is S or NR₅ and that one of X and Y cannot be S at the same time the other of X and Y is NR₅;

(5) Z is H, lower alkyl, aryl, aralkyl, OC(O)R₃ wherein R₃ is independently as defined above, C(O)OR₄ wherein R₄ is independently as defined above, C(O)R₃ wherein R₃ is independently as defined above, CH-(R₁')CO₂R₂' wherein R₁' and R₂' are independently as defined above, halogen, trifluoromethyl,




wherein R, R₁ and R₂ are independently as defined above, heteroaryl, or heteroaralkyl; with the overall proviso that when one of R, R₁, and R₂ is 2-hydroxy, X is O, Y is N and Q is CH=CH, then Z cannot be H or alkyl, and also with the overall proviso that when R, R₁, and R₂ are hydroxy or lower alkyl, Y and X are N or NH, n is zero then Z cannot be furyl or phenyl unsubstituted or substituted with halogen, trifluoromethyl, alkyl, alkoxy or NO₂ which comprises

(1) reacting a compound of the formula




wherein R, R₁, and R₂ are as defined above; and a compound of the formula




wherein $\text{--} = \text{--}$ and Z is as defined above Y and X are O and N, N and O, S and N or N and S, and Ph is phenyl, in the presence of a suitable base; to obtain the compound of formula I wherein Z is as defined above, Y and X are O and N, N and O, S and N, N and S, and R, R₁, and R₂ are as defined above;

(2) treating a compound of the formula




wherein R, R₁, R₂, and Z are as defined above and Q is HC=CH or (CH₂)_n; with hydrazine, and optionally hydrogenating the $\text{--} = \text{--}$ bond when it is a double bond; to obtain a compound of formula I wherein X and Y are N and NH, and R, R₁, R₂, and Z are as defined above;

(3) (step a) treating a compound of the formula



wherein R, R₁, and R₂ are as defined above; with a compound of the formula




wherein Li is lithium, W is H or CO₂R₄ wherein R₄ is as defined above and Z is as defined above; and then (step b) treating the product of the step a with hydrochloric acid in methanol; to obtain a product of formula I wherein X and Y are O and N and Q is $\text{CH} = \text{C} \begin{smallmatrix} \text{---} \\ \text{W} \end{smallmatrix}$,

and R, R₁, R₂, and Z are as defined above;

and, alternatively, further treating the compound obtained in step b with Mo(CO)₆ and then hydrazine to obtain the compound of the formula I wherein X and Y are N and NH and R, R₁, R₂, or Z are as defined above;


(4) treating a compound of the formula



wherein R, R₁, and R₂ are as defined above, with N-chlorosuccinimide and base followed by treating with a compound of the formula



or




to obtain the compound of formula I wherein X is N and Y is O, Q is $\text{HC}\equiv\text{CH}$ and R, R₁, R₂, and Z are as defined above.


(5)a) treating a compound of the formula I of Claim 1 wherein one of X and Y is oxygen and the other is nitrogen with acetonitrile in the presence of molybdenum hexacarbonyl; and then

b) treating with hydrazine to obtain a compound of the formula I of Claim 1 wherein X and Y are nitrogen

(6)a) contacting a compound of the formula




wherein R, R₁ and R₂ are as defined above and a compound of the formula




wherein Z and R₄ are as defined above, and one of X and Y is NH and other is N, or one of X and Y is O and the other is N, and then in situ either

b) dehydrating and decarboxylating the product of step (a) wherein R₄ is hydrogen to obtain a compound of the formula




wherein one of X and Y is NH and the other is N or one of X and Y is O and the other is N, and R, R₁, R₂ and Z are as defined above; or

c) dehydrating the product of step (a) (in situ) wherein R₄ is alkyl to obtain the compound of formula



wherein one of X and Y is NH and the other is N, or one of X and Y is O and the other is N, and R, R₁, R₂ and R₄ are as defined above.

2. A Process according to claim 1 for the preparation of a compound of Claim 1 wherein Q is (CH₂)_n wherein n is zero or two, CH=CH, or



wherein R₄ is as defined above, and wherein X and Y are N and NR₅ with R₅ as defined above, NR₅ and N, O and N or N and O, S and N or N and S.

3. A process according to claim 1 for the preparation of 3-[β-(4'-hydroxy-3'-methoxyphenyl)ethenyl]-5-methylpyrazole or

3-[β-(4'-acetoxy-3'-methoxyphenyl)ethenyl]-5-methylpyrazole

4. A process according to claim 1 for the preparation of 5-[β-(4'-hydroxy-3',-5'-bis(1,1-dimethylethyl)-phenyl)ethenyl]-3-methylisoxazole or

5-[β-(4'-hydroxy-3',5'-dimethoxyphenyl)ethenyl]-3-methylisoxazole.

5. A process according to claim 1 for the preparation of 3-[β-(4'-hydroxy-3',5'-bis(1,1-dimethylethyl)-phenyl)ethenyl]-5-methylpyrazole.

6. A sunscreen composition for use as a sunscreen comprising an ultraviolet absorbing amount of a compound of Claim 1 not including the overall provisos and a suitable carrier therefor.

7. A method of use of a compound of general formula I according to claim 1 alone or in combination with other pharmaceutically active compounds for the manufacture of pharmaceutical compositions for treating inflammation, arthritis, ulcers, allergies, asthma, psoriasis and cardiovascular conditions in mammals suffering therefrom or for the manufacture of compositions for protection from ultraviolet light.



DOCUMENTS CONSIDERED TO BE RELEVANT			EP 87106822.7
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
A	US - A - 4 564 684 (COPP et al.) * Abstract * --	1,10	C 07 D 231/04 C 07 D 231/06 C 07 D 231/12
A	EP - A1 - O 165 782 (ELI LILLY AND COMPANY) * Claims 1,9 * --	1,10	C 07 D 261/02 C 07 D 261/04 C 07 D 261/06 C 07 D 261/08
A	EP - A1 - O 162 383 (BAYER AG) * Abstract * --	1,10	C 07 D 275/02 A 61 K 31/415 A 61 K 31/42
A	US - A - 4 153 707 (MOON) * Abstract * --	1,10	A 61 K 31/425 A 61 K 7/42
D,A	EP - A1 - O 005 192 (BASF) * Abstract * --	1,10	TECHNICAL FIELDS SEARCHED (Int. Cl.4)
A	EP - A2 - O 165 784 (ELI LILLY AND COMPANY) * Claims 1,10 * ----	1,10	C 07 D 231/00 C 07 D 261/00 C 07 D 275/00
The present search report has been drawn up for all claims			
Place of search VIENNA		Date of completion of the search 21-08-1987	Examiner BRUS
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☒ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☒ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

This Page Blank (uspto)